

Index to Volume 14

INDEX OF ORIGINAL ARTICLES

- AB-103. See Benzyl-N-bis(ethylenimido) phosphorocarbamate
- Adenoma, carcinomatosis of, primary in ovaries, attempted palliation by radical surgery in, 384-388
- malignant lymphomas of, in African children, clinical syndrome of, 264
- AMEL, M. R., and GOSLING, J. R. G.: Intracerebral and infiltrative carcinoma of vulva: Bowen's type, 318-329
- See also*, 330-343
- Ad(s). See specific names
- ACTH. See Adrenocorticotrophic hormone
- Actinomycin C (Sanomycin), in combined surgery and chemotherapy in treatment of malignant tumors, use of, 1197-1204
- ADAIR, FRANK E., *jr.* *sw.*, 210-212
- Adamantinoma(s), melanotic, congenital melanocarcinoma, retinal anlage tumor, prognoza, and pigmented epulis of infancy, report of case in adult, 1090-1108
- Adenocarcinoma(s), of ovary and uterus, occurring as coexistent or sequential primary tumors, report of, 817-826
- Adenocarcinoma(s), as type of cancer of lung, validity and value of histological and cytological classifications in relation to, 673-698
- large bowel, treated by 5-fluorouracil (5-FU), clinical evaluation of, 136-137, 139, 140-145, 146
- lung, cytological use of "wet films" in detection of, 1140-1141
- ovarian, treated by 5-fluorouracil (5-FU), clinical evaluation of, 138, 139, 140-145, 150-151
- stomach, treated by 5-fluorouracil (5-FU), clinical evaluation of, 138, 140-145
- Adenoma(s), bronchial, cytological use of "wet films" in detection of, 1142
- Ad(s). See "Apparent diagnostic gap"
- Adrenal(s), malignant lymphomas of, in African children, pathological entity of, 274-275
- Adrenal cortex, response of, in rats inoculated with carcinosarcoma 256 Walker cells, after thermal stress and body temperature changes, study of, 111-116
- width of, in carcinoma of lung, report on, 105-110
- Adrenalectomy, and cortisone therapy, effect of, on advanced breast carcinoma, evaluation of, 1259-1265
- in "recurrent" or inoperable breast carcinoma, results of, 304-307
- associated hormonal therapy and, 304-306
- relationship to stage of disease in, 306, 307
- Adrenal medulla, ganglioneuroma of, containing cells morphologically identical to hilus cells, report of, 421-425
- case report, 421
- criteria necessary for identification of, 423-425
- crystalloids of Reinke in, 421, 422, 423-425
- ganglion cells in, 421, 422, 423-425
- interpretations of origin of cells in, 423-425
- gonadal and juxtagonadal location of, 424-425
- possible protein nature of, 424
- microscopic appearance of tumor in, 421-422
- mild chronic cholecystitis associated with, 421
- unique nature of tumor in, 423
- use of Masson's polychrome stain in, 421, 422
- Adrenocorticotrophic hormone (ACTH), effect of, on transplanted tumor in mice, report of, 507-511
- for giant follicular lymphoblastoma, 771, 772-773
- Adrenal steroid(s), role of, in development of gastroduodenal metastases from breast carcinoma, 426-431
- Age, as grouping factor in measurement study of cells exfoliated from female genital tract, report of, 442
- "Aging," of carcinosarcoma 256 Walker cells, effect of, on development of tumors in rats, report of, 1057-1059
- AGOSTINO, D., *jr.* *sw.*, 957-962
- Alcohol, as etiological factor in cancer of esophagus, study of, 389-413
- ALD. See Aldolase
- Aldolase (ALD), levels of, in serum of patients with carcinoma of breast, report of, 726-733
- Alimentary tract. See also specific names of organs
- involvement of, by multiple lymphomatous polyposis, 247-256
- Alkaline phosphatase(s), quantitative assessment of activity of, in growth and differentiation patterns of normal and malignant cells, report of, 502-506
- AMBRUS, JULIAN L., *jr.* *sw.*, 853-861
- Ameloblastoma(s), as epithelial odontogenic tumors in man, classification, histopathology, and clinical behavior of, 76-80
- comparison of, with ameloblastic adenomatoid tumors, 76, 81
- Amethopterin, use of, in prolonged intra-arterial chemotherapy technique for treatment of inoperable cancer, 493-495
- 4-Amino-N¹⁰-methyl pteroylglutamic acid (amethopterin, Methotrexate), continuous intra-arterial administration of, with intermittent intramuscular citrovorum factor (CF, Leucovorin) antimetabolite-metabolite cancer chemotherapy in 744-752
- carotid arterial catheterization method in, 744-745
- case reports, 749-750
- catheterizing of internal and common carotid arteries in, 746
- dose schedules used in, 747-748
- formation of clots in, 751
- hypogastric artery catheterization method in, 746-747
- incidence of generalized infection in, 750
- of leakage in 750-751
- of local infection in, 750
- infusion assembly in, 747
- possible more efficient methods of catheterization in, 751
- preventive care of pump in, 751
- sites of arterial catheterization and incidence of complications of technique in, 748-752
- toxicity in infusion area in, 751
- use of fluorescein (Fluorescein) in, 747
- in study of growth curves of human cancer after transplantation, report of, 1117-1121
- in treatment of malignant lymphomas in African children, use of, 266
- 6-Aminonicotinamide (6-AN), in therapy of disseminated cancer, clinical use of, 644-648
- Amputation, intercapulothoracic, as treatment of postmastectomy lymphangiosarcoma, effect of, 210-212
- Amylase(s), activity of, in body fluids, study of levels of, 1109-1114
- 6-AN. See 6-Aminonicotinamide
- Anatomy, of larynx, in relation to dissemination of cancer, 1131-1138
- ANDERSON, PHILIP C., *jr.* *sw.*, 102-104
- Androgen(s). See specific names
- Androstenediol dipropionate (5 α -androstan-3 α ,17 β -diol), in treatment of breast cancer, effect of, on clinical course and hormonal excretion, 707-710
- Androsterone, effect of, on clinical course and hormonal excretion in cancer of breast, report of, 195-198
- Angiosarcoma(s), in children, as part of malignant compound tumor, 524-533
- ANGLEM, THOMAS J., and GRAY, EDWARD B., *Jr.*: Cancer of the breast and biological predetermination: a re-evaluation, 1122-1126
- Anoxia, as physical agent affecting radiation sensitivity of mouse tumors, report of, 122-123
- Anus. See also Colon and Rectum
- carcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 242
- "Apparent diagnostic gap" (ADG), in biostatistical approach to tumor growth, report of, 1272-1294
- "Apparent survival time" (AST), in biostatistical approach to tumor growth, report of, 1272-1294
- Argentaffin(s), in adenomatoid bronchial tumors, presence of, 171-173
- Argyrophil(s), in adenomatoid bronchial tumors, presence of, 171-173
- Arm(s), upper, lymphedematous, post-mastectomy, treatment of lymphangiosarcoma developing in, 210-212
- ARONS, MARVIN S.; KETCHAM, ALFRED S., and MANTEL, NATHAN: The effect of growth hormone and ACTH on a transplanted tumor, 507-511
- SMITH, ROBERT R., and MYERS, MAX H.: Significance of cancer cells in operative wounds, 1041-1044
- Arsenic, as ingested carcinogenic agent in Bowen's type of intracerebral and infiltrative carcinoma of vulva, possible role of, 321-328
- Artery(ies), continuous administration of intra-arterial 4-amino-N¹⁰-methyl pteroylglutamic acid (Methotrexate) via, with intermittent intramuscular citrovorum factor (CF), in antimetabolite-metabolite cancer chemotherapy, report of, 744-752
- Arthritis, skeletal lesions of acute leukemia and, 616
- AST. See "Apparent survival time"
- Atelactasia, postoperative, use of pancreatic deoxyribonuclease (dornase) in treatment of, 415-416, 420
- segmental, postoperative, use of pancreatic deoxyribonuclease (dornase) in treatment of 415-416, 420
- Atypia(s), cell, in cervical biopsy specimens, value of serial step sectioning after finding, 213-216
- Autotransplantation, of human cancer, quantitative studies of, 971-978
- histology of original and transplanted tumor tissues in, 972-975
- injections of subcutaneous homotransplants of tissue-cultured human cancer cells of H.Ep-2 and/or H.Ep-3 cell lines in addition to, in, 972
- preparation of cancer tissue obtained at laparotomy or biopsy in, 971
- preliminary cell counts on, 971
- second cell counts on, 971
- use of Gey's salt solution in preparation of, 971
- of Giemsa-stained smear from most concentration suspensions in, 972
- procedure for tumor cell suspension injection used in, 972
- sources of error in evaluating results in, 975-978
- study of auto- and homo-transplant growth in, 975
- summary of experience in, 972-978
- amount of cells in inoculation and number of patients with growth in, 972
- time of appearance of palpable autotransplants in, 972-977
- 8-Azaguanine (Aza), in combined surgery and chemotherapy as treatment for malignant tumors, use of 1197-1204
- Aza (8-Azaguanine), in combined sur-

- gery and chemotherapy as treatment for malignant tumors, use of, 1197-1204
- BAGGENSTOSS, ARCHIE H., *jt. su.*, 221-230, 231-237, 238-248, 901-912
- Baker interference microscope, in quantitative assessment of alkaline phosphatase activity in growth and differentiation patterns of normal and malignant cells, use of, 502-506
- BARDOS, THOMAS J., *jt. su.*, 853-861
- BARTLETT, RAYMOND C.; OTIS, RICHARD D., and LAAKSO, ANDREW O.: Multiple congenital neoplasms of soft tissues. Report of 4 cases in 1 family, 913-920
- BAUER, WALTER C., *jt. su.*, 55-66
- BECKER, STANLEY M., *jt. su.*, 1235-1250
- BECKETT, V. L., *jt. su.*, 131-152
- BELOVICH, DORIS, *jt. su.*, 1139-1143
- Benecarbimide. *See* Benzyl-N-bis(ethylenimido)phosphorocarbamate
- Benzo[a]pyrene (BP), as carcinogenic chemical, in newborn mice, report of, 308-317
- role of, in carcinogenesis, experimental study of, 1306-1315
- Benzyl-N-bis(ethylenimido)phosphorocarbamate, (benecarbimide, AB-103), as dual antagonist in treatment of bronchogenic carcinoma, use of, 853-861
- BERG, JOHN W., *jt. su.*, 539-546
- BESSE, BYRON E., JR., *jt. su.*, 608-621
- BHATTACHARYA, K., *jt. su.*, 502-506
- Biostatistics, in study of tumor growth, use of, 1272-1294
- Biometrics, in study of etiology of cancer of cervix, report of, 153-161
- Biopsy, cone, microscopic study of specimen from, predictability of residual carcinoma in uterus after treatment for intraepithelial carcinoma of cervix, report of, 795-800
- 2,5-Bis(ethylenimino)-3,6-dipropoxy-1,4-benzoquinone (E-39), in study of combined surgery and chemotherapy in treatment of malignant tumors, use of, 1197-1204
- Bladder, epitheliomas of, as multiple primary malignant tumors of multicentric origin, report of, 241
- BLAND, J. E.; WINDMILLER, JOAN; WHITAKER, JOANNE, and CLARK, KEMP: Intrathecal cyclophosphamide (Cytosan) in dogs, 1115-1116
- BLOCK, GEORGE E., *jt. su.*, 1081-1084
- Blood. *See also* Serum (a)
- carcinosarcoma 256 Walker cells in, effect of human fibrinolysin on survival of, 957-962
- of cancer patients, excretion and distribution studies of radioactive-labeled alkylating drugs in, 953-956
- serum, human, of normal individuals and individuals with various pathological conditions, triosephosphate isomerase activity of, 1166-1174
- BOND, VICTOR P., *jt. su.*, 639-643
- Bone(s), endochondral, alteration in formation of, in cases of acute leukemia, report of, 618-619
- involvement of, by giant follicular lymphoblastoma, report of, 762-763
- in cases of adenoid cystic carcinoma, report of, 1241-1250
- jaw, malignant lymphomas of, in African children, pathological entity of, 270-272, 273
- lesions, in acute leukemia, 608-621
- sarcoma of, complicating Paget's disease, report of, 517-523
- age of patient at time of diagnosis of sarcoma in, 519, 520-521, 523
- case reports, 517-520
- histological type of sarcoma in, 521
- pathological diagnosis in, 519
- physical findings in, 517, 519, 523
- prognosis in, 523
- roentgenographic findings in, 517-518, 520, 521, 522
- site of Paget's disease in, 517-523
- of sarcoma in, 517-523
- surgical procedures used in, 517, 519, 520, 521
- survival time after diagnosis of
- Bone(s), sarcoma of—(Cont'd.)
- sarcoma in, 517-523
- BOSTROM, R. C., *jt. su.*, 437-454, 455-468
- Bowen's disease, of vulva, differentiating features of, 318-329
- differentiation between Paget's disease of vulva and, 318, 329
- occurrence of other malignant neoplasms and, 320-321, 328-329
- BOWERS, C. Y., *jt. su.*, 195-198
- Br (benzo[a]pyrene), as carcinogenic chemical, in newborn mice, report of, 308-317
- BRADY, LUTHER W., *jt. su.*, 1235-1250
- Brain, cell-free filtrates from leukemic mature AKR mice, failure to demonstrate acceleration of leukemia in animals of same strain, report of, 496-501
- giant cell fibrosarcoma of, 841-852
- anatomical location of tumors in, 842-845
- average survival of patients in, 842-845
- differentiation from malignant astrocytoma in, 851
- extent of tumor involvement in, 842-845
- histological findings in, 842-850
- appearance of cytoplasm in, 846-847
- of nuclei in, 846-847
- bizarre giant cells in, 847-852
- fibrous element in, 847, 849
- mitoses in, 847-851
- presence of anaplasia in, 845-846, 849
- reticular reaction in, 847, 849-850
- transition forms in, 847, 849
- use of hematoxylin and eosin stain in, 845
- vascularity in, 849-850
- lack of neuronal and glial elements in, 852
- metastases, value of radiation therapy in management of, 577-581
- age distribution of patients with, 577
- confirmation of clinical diagnosis in, 577
- distribution according to site of primary tumor in, 577, 578
- possibility of selection in, 580-581
- results of, 578, 579-581
- dose-response correlation in, 580-581
- remissions after, 579-581
- response to first and second courses of irradiation in, 579-580
- survival rates in, 579-581
- symptoms most frequently found in patients with, 577
- type of irradiation in, 577-579
- BRANDBOG, LLOYD L.; TANIGUCHI, LILIAN, and RUBIN, CYRUS E.: Is exfoliative cytology practical for more general use in the diagnosis of gastric cancer? A simplified chymotrypsin technique, 1074-1080
- Breast, cancer of, advanced, therapeutic oophorectomy for, significance of metastases to ovary and ovarian cortical stromal hyperplasia in, 126-130
- classification of cases in, 127
- metastases to ovaries and, 127-128
- results for other types of therapy used in, 128
- survival period after breast operation in, 128
- survival period after oophorectomy in, 127-128
- ovarian cortical stromal hyperplasia and, 128, 129-130
- survival after breast operation in, 128, 129
- survival after oophorectomy in, 128, 129
- and biological predeterminism, re-evaluation of, 1122-1126
- changes in descriptions of tumors in, 1122
- comparison of "early" and "late" series in, 1122
- operative procedures and indications for surgery in, 1122
- relation of duration of symptoms to 5-year survival rates in, 1123
- Breast, cancer of, advanced—(Cont'd.)
- statistical analyses of rates recorded in, 1123-1125
- possible "distortion" factors in, 1123-1125
- survival rates in, 1122-1123
- metastatic involvement and, 1122
- ulceration of tumors and, 1122
- effect of androstenediol dipropionate (5 α -androstane-3 α ,17 β -diol) on clinical course and hormone excretion in, 707-710
- administration of compound in, 707
- criteria for recording objective response to therapy in, 708
- incidence of virilization in, 709
- luteinizing hormone levels in, 708-709
- method of studying hormonal balance in, 708
- P and S urine levels in, 708-709
- randomization procedures, 707
- type of dominant lesions in, 709
- urinary hormonal excretion studies in, 708-709
- creatinine levels in, 708-709
- creatinine levels in, 708-709
- fluorometric phenol levels in, 708-709
- formaldehydogenic corticoid levels in, 708-709
- gonad-stimulating hormone levels in, 708-709
- 17-ketosteroid (17-Ks) levels in, 708-709
- pragnadiol chromogen levels in, 708-709
- prolactin levels in, 708-709
- urine calcium in, 708-709
- metastatic, effect of androsterone on clinical course and hormonal excretion in, 195-198
- categories of patients, by menopausal status, in, 195
- subdivision by systemic involvement in, 195
- distribution of patients and response to therapy in, 195, 196
- dose schedule used in, 196
- randomization schema applied in, 195-196
- remissions after, 196-198
- side effects after, 197
- treatment protocol in, 195
- urinary excretion levels before and during administration in, 196-197
- 17-ketosteroid (17-Ks) levels after therapy with, 196, 197, 198
- significance of occult metastases in, 1266-1271
- location of regional lymph nodes in, 1266
- mammary specimens used in study of, 1266
- method of obtaining 5-year survival data in, 1266
- other clinical factors involved in prognosis in, 1267-1271
- anatomical factors as, 1268, 1269
- location of tumor as, 1268, 1270
- patient age as, 1268
- size of primary tumor as, 1268-1269
- surgical procedure as, 1268
- type of carcinoma as, 1268, 1269
- proposed method for routine sectioning in, 1270-1271
- selection of specimens for serial sectioning in, 1266
- significance of lymph node involvement on survival rates in, 1266-1271
- number of involved axillary nodes and, 1267
- grid analysis used in, 1267
- selective nature of patient group studied in, 1267
- some factors causing patient delay in seeking treatment for, 979-988
- carcinoma of, advanced, comparative evaluation of adrenalectomy and cortisone in treatment of, 1259-1265

Breast, carcinoma of, advanced—(Contd.)

- case reports, 1261-1262
criteria for remission in 1260
follow-up roentgenograms in, 1260
method of administering cortisone acetate in, 1259
dose schedule used in, 1259
randomization procedure used in, 1259
response to adrenalectomy in, 1260-1261
effectiveness of adrenalectomy after failure to respond to cortisone therapy in, 1260
ovarian function and response to adrenalectomy in, 1260
response to corticosteroid therapy in, 1260
routine check-ups in, 1260
site of metastases and response in, 1260-1264
survival period postadrenalectomy in, 1260-1264
bilateral, as multiple primary malignant tumors of multicentric origin, report of, 242-243
duct epithelium, and fibroadenoma, method of labeling tissues with tritiated thymidine in vitro and its use in comparing rates of cell proliferation in, 639-643
comparison of deoxyribonucleic acid (DNA) in 3 cells populations in, 639-643
human tissue studies in, 640-643
calculated rates of proliferation in, 642
duration of DNA synthesis in, 642
equations for generation times in, 642
method of labeling cells in, 641
results of cell counts and generation times for tissues studied in, 639, 642-643
preliminary studies in, 639
chief difficulty encountered in labeling cells in, 640
transplantable mouse sarcoma used in, 639
use of BNL albino mice in, 639
female, metastatic hormonal therapy with 2 α -methyl dihydrotestosterone propionate in, 705-706
criteria for objection remission in, 705
dose of hormone used in, 705
double-blind technique of drug administration in, 705
incidence of virilizing effects in, 706
percentage remissions in, 706
structure of compound in, 705
types of virilizing effects in, 706
gastroduodenal metastases from, adrenal steroid-induced, 426-431
autopsy material used in, 426
average duration of adrenal steroid therapy in, 427
patient age at death in, 427
clinical significance of metastatic sites in, 429-430
effect on local tissue resistance in, 429
incidence of ulcerations of stomach and duodenum in, 427-428
investigation of hormones, drugs, and surgical procedures used in patients included in study of, 426
previous studies with cortisone in, 428-429
previously reported rates of, 426
sites of metastases in, 427-430
types of metastases in, 427-430
gelatinous, prognostic significance of, 699-704
age distribution and 5-year survival rates, by gelatinous categories, in, 700, 701
clinical presentation of tumors in, 700
correlation of survival rates

Breast, carcinoma of, gelatinous—

- (Contd.)
with pathological findings in, 700, 701-703
grading of tumor in, 703
influence of axillary metastases in, 702
of tumor size in, 702
location of primary tumor in, 703-704
multiple primary tumors in, 704
recurrence after operation in, 704
site of first metastases in, 704
follow-up data in, 700
gross appearance of tumor in, 700
incidence of axillary metastases in, 700
microscopic appearance of tumor in, 700-701
surgical specimens in, 700
tumor categories in, 700-704
isocitric dehydrogenase, malic dehydrogenase, and glycolytic enzyme levels in patients with, 726-733
formula for calculating correlation coefficients in, 728
incidence and degree of elevations of enzymes in, 728-732
method of determining enzyme levels in serum in, 727-728
patient categories in, 726, 727
summary of correlations for different enzymes in, 731
types of therapy in, 726, 727
upper limits of normal values for serum enzymes in, 727
male, development of, after estrogen treatment for prostatic cancer, report of, 234
metastatic, alterations in serum proteins of patients receiving palliative therapy for 1081-1084
comparison of electrophoretic and chemical separation in, 1083
concentration of serum protein fractions found in, 1082, 1083
fractionation method used in, 1081
moving boundary or paper electrophoresis used in, 1081
organs involved in, 1082-1083
samples of clotted blood used in study of, 1081
types of treatment used in, 1081
primary, frequency of occurrence of second primary malignant tumors in women with, 232
"recurrent" or inoperable, effect of hormonal treatment of, and survival of patients with, 300-307
age distribution and survival time in, 300-301
comparison of patient survival with and without hormonal therapy in, 302, 303
definition of "recurrent" in, 300
evaluation of objective and subjective improvement in, 302, 304
forms of therapy in, 302-303
palliative effects of, 302-307
relationship of adrenalectomy and 303-304, 305, 306
results of different kinds of hormonal therapy in, 303, 305
time of and type of recurrence in, 301-302, 303
location of metastases and, 301, 302, 303
survival time and, 301-302
types of lesions improved by hormonal therapy in, 302, 304
5-year survival rates in, 306
treatment by 5-fluorouracil (5-FU), clinical evaluation of, 137-138, 140-145, 146-147, 148
mass in, some observations about patients with, 979-988
comparison of apparently pertinent factors between cancer and non-cancer patients and those patients who do or do not delay in, 987

Breast, mass—(Contd.)

- data for patients with clinically benign lesions not operated upon in, 980, 981
definition of delay in, 981
delay time for patients with and without cancer in, 984-985, 987
comparison of, with previously reported delay times in, 984, 987
effect of public education about cancer on patients in, 985
factors investigated in, 982-987
follow-up of patients in, 981
incidence of breast cancer in, 981
psychological "set" of patients about breast cancer in, 983-987
types of emotional and delay factors previously recorded in, 979-980
BRENNAN, M. J., JR. SM., 131-152
BROGHAMER, WALTER L., JR., and CHRISTOPHERSON, WILLIAM M.: An interferometric study of the anhydrous nuclear mass of exfoliated cells from experimental cervical cancer, 378-383
Jt. SM., 201-204
Bronchoscopies, use of pancreatic deoxyribonuclease (dornase) prior to, 419, 420
use of "wet film" study at time of, in detection of lung carcinoma, report of, 1139-1143
Bronchus(i), and trachea, of smokers and nonsmokers, microscopic properties of basement membrane and elastic fibers of, 1175-1182
carcinoids and cylindromas of, non-benign nature of, 539-546
clinical presentation in, 539-540
course of disease and results of treatment for bronchial carcinoids in, 544-545
incidence of metastases in, 545
irradiation therapy in, 544-545
surgery in, 544-545
5-year survival rates in, 545
course of disease and results of treatment for cylindromas in, 545-546
irradiation in, 545-546
surgery in, 545-546
survival rates in, 545-546
microscopic findings in, 542-544
correlation between degree of nuclear anaplasia and incidence of metastases in, 542, 543-544
glandular differentiation in, 542
marked pleomorphism in, 543
mucus secretion in, 542
pathological findings in, 540-544
extrabronchial extension in, 542
grading of carcinoid tumors in, 542, 543
gross description of tumors in, 540-542
location of tumors in, 539
types of patients with, 539
consideration of carcinoids and salivary tumors of, 161-178
case reports, 162-167, 167-169
classification of tumors in, 175
clinical features of carcinoid syndrome in, 170-171, 173
comparison of carcinoid and adenoid cystic types in, 174
hamartomas in, 169
histogenesis of adenoid cystic carcinomas in bronchus in, 174
histological findings for carcinoid group in, 165-167
cells in, 163-167
over-all pattern in, 163-167
stroma in, 163-167
histological findings for salivary gland type tumor group in, 167-169
cylindromatous pattern in, 167-169
mucus-secreting and epidermoid pattern in, 168, 169
Kulschitzky cell distribution in, 172
"mixed" tumors with carcinoid and adenoid cystic features in, 174
origin, in relation to bronchial epithelium, of, 173
patient age in, 170
sex in, 170

- Bronchus (i), carcinoids and salivary tumors of—(Contd.)
 presence of mucin in, 172
 role of 5-hydroxytryptamine (5-HT) in, 171
 of metastases in, 170
 serotonin in, 172, 173
 staining reactions of carcinoids in, 171-173
 argentaffin reaction in, 171-173
 argyrophilic reaction in, 171-173
 silver techniques in, 171
 tumors similar to cylindromas in, 174-175
 urinary 5-hydroxyindoleacetic acid (5-HIAA), levels in, 172, 173
 carcinoma of, use of benzyl-N-bis(ethylenimido)phosphorocarbamate (AB-103) in treatment of, 853-861
 chemical structure of, 853
 comparison of, with ethyl-N-bis(ethylenimido)phosphorocarbamate (AB-100), in, 853
 dosage given in, 856
 regression of soft tissue metastases in, 856, 860-861
 survival rates in, 856, 859-861
 therapeutic results in, 856, 858-861
 objective response after, 856, 858-861
 subjective improvement after, 856, 858-861
 toxicity in, 856-858
 use of weekly maintenance doses in, 861
 tree, dysplastic lesions of, 1226-1234
 clinical data for men without lung cancer, arranged in order of increasing age, in, 1227
 for patients with lung cancer, in 1229-1231
 for women without lung cancer, arranged in order of increasing age, in, 1228
 comparison of extent of metaplasia in men between nonsmokers and medium to heavy smokers in, 1230, 1231
 of mucosal hyperplasia in different smoking categories in, 1230
 in men and women "paired" for age and smoking history in, 1230
 of thickness of mucosa and degree of metaplasia in male smokers with and without lung carcinoma in, 1230, 1231
 diagram of bronchial segments that were studied in, 1227
 fixation of sections used in, 1228
 focus of present investigation of, 1227-1228
 grading of metaplasia in, 1228
 of smoking categories in, 1229
 measurement of mucosa in, 1228
 preparation of cytological specimens in, 1229
 previous investigations of histogenesis of, 1226-1227
 staining of sections in, 1228
 use of routine postmortem material and surgically removed lungs in, 1127
 BROOKS, VIVIAN, *ft. sm.*, 1259-1265
 BROSS, IRWIN J., *ft. sm.*, 389-413
 BRUNSCHWIG, ALEXANDER: Attempted palliation by radical surgery for pelvic and abdominal carcinomatosis primary in the ovaries, 384-388
ft. sm., 582-586, 971-978
 BULLOCK, WELDON K., *ft. sm.*, 512-516
 BURKITT, DENIS, and O'CONNOR, G. T.: Malignant lymphoma in African children. I. A clinical syndrome, 258-269
 BUTCHER, HARVEY R., *ft. sm.*, 1251-1258
 Calcium, renal, excretion, in advanced cancer, effect of citrate infusion on, 934-943
 CAMPBELL, J. S.; MAGNER, D., and FOURNIER, PIERRE: Adenocarcinomas of ovary and uterus occurring as co-existent or sequential primary neoplasms, 817-826
 Cancer(s). *See also* Carcinoma(s) and specific anatomical sites and types of cancer
 abdominal, advanced metastatic, primary in ovaries, attempted palliation by radical surgery of, 384-388
 advanced, hypercitricemia and parathyroid function in, 934-943
 clinical data for patients receiving infusions in, 935
 comparison of excretion levels with those after saline infusion in, 940
 correlation between spontaneous hypercalciuria and hypercitricemia in, 939-940
 division of cancer patients into groups in, 936
 effect of infusion upon estimated serum ionized calcium in, 936-937
 of sodium chloride infusion on sodium citrate in, 938
 healthy volunteers and patients included in study of, 934
 infusion of sodium citrate in, 934, 942
 method of estimating ionic calcium in, 935-936
 overnight calcium excretions and serum citrate and calcium concentration prior to infusion in, 938
 possibility of level of citrate in blood modifying hormone secretion in, 941
 significance of, 941
 preliminary urine collections in, 934
 previous reports of hypercitricemia in cancer patients, 934, 935
 renal citrate and calcium excretion after infusion in, 938
 response of serum calcium and phosphorus concentration to citrate infusion in, 940
 urinary calcium excretion after infusion in, 937-939
 phosphate excretion after infusion in, 937
 response to citrate infusion in hypercitricemic and hypercalcemic patients in, 940
 quantitative studies of autotransplantation in, 971-978
 antimetabolite-metabolite chemotherapy of, using continuous intrarterial 4-amino-N¹⁰-methyl pteroylglutamic acid (Methotrexate) with intermittent intramuscular citrovorum factor (cv), 744-752
 breast, advanced, therapeutic oophorectomy for, significance of metastases to ovary and of ovarian cortical stromal hyperplasia in, 126-130
 and biological predeterminism, re-evaluation of, 1122-1126
 statistical analyses of rates recorded in, 1123-1125
 possible "distortion" factors in, 1123-1125
 effect of androstanediol dipropionate (5 α -androstan-3 α ,17 β -diol) therapy on clinical course and hormonal excretion in, 707-710
 metastatic, effect of androsterone on clinical course and hormonal excretion in, 195-198
 significance of occult metastases in, 1266-1271
 cells, in operative wounds, significance of, 1041-1044
 cervical, biometric approach to etiology of, 153-161
 critical study of biopsy in diagnosis of, 213-216
 experimental, interferometric study of anhydrous mass of exfoliated cells from, 378-383
 treated by radiation, development of in situ epidermoid carcinoma of cervix and vagina after, 353-360
 chemotherapy of, with 5-fluorouracil (5-FU), clinical evaluation of, 131-152
 use of naphthylchloroacetamide in, 296-299
 disseminated, clinical use of 6-aminonicotinamide (6-A), in patients with, 644-648
 Cancer(s), disseminated—(Contd.)
 administration of 6-AN in, 644
 antineoplastic potency of 6-AN in, 644-647
 case reports, 645-646
 clinical evaluation of results of, 646
 clinical toxicity in, 647
 dose of 6-AN used in, 644, 646
 duration of therapy in, 645-647
 effects of 6-AN therapy on in vitro incorporation of formate-C¹⁴ into tumor proteins in, 644, 645, 646, 647
 types of tumors in, 644, 645
 esophageal, study of etiological factors in, 389-413
 excretion and distribution studies of radioisotope-labeled alkylating drugs in, 953-956
 generalized, with and without liver metastases, activity of triphosphate isomerase in blood serum from, 1066-1074
 human, transplants of, effect of amethopterin (Methotrexate) on growth curves of, report of, 1117-1121
 in Singaporean children, epidemiological study of, 534-538
 cancer death rates in, 536
 distribution of types of cancer in, 535
 comparison with rates reported from other countries in, 535-536
 geographical location of Singapore in relation to, 534
 infant mortality rate in relation to, 534
 leukemia incidence in, 535, 536-537
 multiracial population in relation to, 534
 population age in relation to, 534
 sources of bias in, 534-535
 age groupings and mean age at death as, 535
 relative sex and age incidence as, 535
 use of postmortem protocols as, 534
 inoperable, prolonged intra-arterial chemotherapy technique for, 493-495
 criteria for pump used in, 494
 design of pump used in, 493, 494
 infusion of amethopterin in, 494
 primary hazards involved in, 495
 laryngeal, anatomical studies related to dissemination of, 1131-1148
 liver, middle hepatic lobectomy for, 1295-1300
 teratoid, in infancy, complete right hepatic lobectomy for, 361-368
 localized, activity of triphosphate isomerase in blood serum from patients with, 1166-1174
 lung, cytological use of "wet films" in detection of, 1139-1143
 peripheral, arising in scars, report of, 1205-1215
 relationship of certain factors to survival of patients treated by pulmonary resections for, 1251-1258
 validity and value of histological and cytological classification of, 673-698
 multiple, primary, of different tissues or organs, 231-237
 hereditary influences in patients with, 234-235
 limitations in study of, 234
 importance of age at time of diagnosis in, 236
 induction of second primary cancer by treatment of initial primary cancer in, 232-234
 estrogen treatment of male prostate cancer and development of male breast cancer in, 234
 role of Stewart-Treves syndrome in, 233-234
 of therapeutic ionizing radiation in, 233
 occurrence rates for family members in, 235
 by age groups, 235
 relationship between cancer of

- Cancer(s), multiple, primary—(Contd.)**
 estrogen-sensitive organ and carcinogenesis in other estrogen-sensitive organs in, 231-232
 of multicentric origin, report of, 238-248
 adenocarcinomas of colon and rectum as, 240
 frequent occurrence of, 240
 multiple polyposis and chronic ulcerative colitis associated with, 240
 simultaneously appearing and interval lesions in, 240
 of stomach as, 239
 possibility of supposed recurrences being new independent neoplasms in, 239
 bilateral carcinomas of breast as, 242-243
 rate of occurrence for interval lesions in, 242-243
 for simultaneous lesions in, 242
 ovarian cancers as, 243-244
 difficulties in distinguishing metastases from true second primary tumor in, 243-244
 panhysterectomy as possible preventive therapy in, 244
 testicular cancers as, 243
 high incidence rate of, 243
 prophylactic orchiectomy as possible preventive therapy in, 243
 carcinoid tumors of small intestine as, 239-240
 incidental finding at necropsy of, 240
 carcinomas of cervix, vagina, vulva, and anus as, 242
 of oral cavity as, 238-239
 associated leukoplakia and, 239
 use of tobacco in patients with, 239
 of parenchymatous organs as, 244
 kidney as site of, 244
 liver as site of, 244
 thyroid and pancreas as site of, 244
 difficulties in differential diagnosis of metastasis and true second primary cancer in, 238-248
 reported versus true incidence of, 238
 epitheliomas of larynx, pharynx, and esophagus as, 239
 associated squamous cell lesions of contiguous epithelium and, 239
 of lips as, 239
 of skin as, 238
 of urinary tract as, 240-242
 bladder as site of, 241
 frequency of, 240
 possible mechanisms of, 240-241
 lesions associated with those of renal pelvis as, 241-242
 gliomas of central nervous system as, 244
 importance of theory of multicentricity in relation to treatment in, 246
 malignant lesions of reticulo-endothelial system as, 244-245
 lack of knowledge on nature and pathogenesis of, 245
 present knowledge about carcinogenesis in relation to, 245-246
over-all report of, 221-230
 age of patients at time of diagnosis in, 221, 229
 by incidence of simultaneous or interval lesions in, 221, 229
 classification of, 226-227
 criteria for diagnosis of, 222-223, 226
 increase in incidence of, 221
 rate of occurrence in, 222, 223
- Cancer(s), over-all—(Contd.)**
 224-225, 227, 228, 229-230
 definition of "epithelioma" in, 230
 exclusion of multiple epitheliomas confined to skin alone in, 228-229, 230
 for multicentric lesions, 223, 224-225, 227, 228, 229-230
 of bilaterally paired organs, 223
 of common contiguous tissue shared by different organs, 223
 of different tissues or organs, 224-225, 227
 of same tissue or organ, 223
 for multicentric lesions plus lesion(s) of different tissues or organs, 227
 for multiple lesions of different organs or tissues, 222, 229
 of childhood, use of chlorambucil in treatment of, 36-41
 case reports, 37-39
 chemical structure of, 36
 data for children in, 36
 dosage regimen used in, 39-40
 objective remissions following drug administration in, 37-39
 relationship to toxic effects of, 39-40
 subjective improvement accompanying drug administration in, 37
 therapeutic effects of, 36-37, 40
 toxic effects after administration of, 39
 types of cancer in children studied in, 36, 37
 of maintenance therapy given in, 40
 ovarian, metastatic, luteinization of stroma of, endocrine significance of, 827-840
 "pedigree," in studies of inheritance of cancer of human uterine cervix, use of, 180, 181
 pelvic, advanced, metastatic, primary in ovaries, attempted palliation by radical surgery of, 384-388
 skin, incidence of, in families of cervical cancer patients, report of, 186-187, 192, 193
 stomach, in spouses of patients with, incidence of, 199-200
 simplified chymotrypsin technique for exfoliative cytology in, 1074-1080
 terminal, psychological problems in management of, 1063-1073
 case reports, 1066-1072
 factor of denial in, 1064-1065
 conscious denial as, 1064
 unconscious denial as, 1064-1065
 factor of isolation in, 1066
 factor of loss of control and mastery in, 1064
 attempts at "magical" solutions in, 1064
 principles of management in, 1072
 reaction to grief in, 1065
 sense of failure in, 1065-1066
 uterine cervix, spread of, as seen in giant histological sections, report of, 42-54
 studies on inheritance of, 179-184
 vulvar, report of, 13-20
Candida, skin test reactions to, in cases of Hodgkin's disease and lymphosarcoma, report of, 597-607
Carcinogen(s). See also specific names
 arsenic as, in Bowen's type of intraepithelial and infiltrative carcinoma of vulva, possible role of, 321, 328
 chemicals as, effects of, on newborn mice, report of, 308-317
 cigarette tar and beta radiation as, in mice, action of, 344-352
 in development of benign epithelial tumors, role of hair follicle as location in, 801-816
 methylcholanthrene as, development of carcinoma of gallbladder in dogs after implantation of, 1127-1130
 in experimental tumors of peripheral nerve trunk in rabbits, re-
- Carcinogen(s), methylcholanthrene as—(Contd.)**
 port of, 1030-1035
 in study of progression of cervical dysplasia in mouse, report of, 201-204
Carcinogenesis, in relation to study of multiple primary malignant tumors of multicentric origin, report of, 238-248
 of benign epithelial tumors, role of hair follicle in, 801-816
 role of acidic fractions of tobacco in, 1306-1315
Carcinoid(s), bronchial, nonbenign nature of, 539-546
 developing in dermoid cysts of ovary, report of, 990, 993-994, 995-999
 gross appearance of, 997
 incidence of carcinoid syndrome in, 997-998
 microscopic appearance in, 997-998
 report of, 161-178
 symptoms in, 998
 treatment for, 997, 998
Carcinoid syndrome, clinical features of, in tumors of bronchus, report of, 170-171, 173
Carcinoma(s). See also Cancer(s) and specific anatomical sites of types of carcinoma
 adenoid cystic, bronchial, report of, 161-178
 clinicopathological study of, 1235-1250
 case reports, 1241-1245
 patient age in, 1237
 race in, 1237
 sex in, 1237
 data on clinical and pathological findings for patients in, 1242-1243
 duration of symptoms at time of diagnosis in, 1245
 interval between first symptom and appearance of visceral metastases in, 1246
 onset of symptoms in, 1245
 period of time from primary treatment of tumor to time of local recurrence in, 1246
 total duration of tumor existence in, 1245
 type of symptoms in, 1245-1246
 differential diagnosis in, 1238-1241
 gross appearance of tumor in, 1237-1238
 high incidence of bone invasion in, 1246
 histogenesis of, 1236
 incidence of metastasis in, 1246-1247
 microscopic features of, 1238
 prognosis in, 1248-1250
 results of treatment for, 1247-1250
 local excision as, 1247
 neurosurgery as, 1248
 radical excision as, 1248
 radiotherapy as, 1247-1248
 results of diagnosis of tumors in, 1236
 site of origin in, 1237
 difficulty in demarcating extent of tumor at time of surgery in, 1238, 1249
 size in, 1237
 slow growth rate in, 1246
 basal cell, metastasizing, report of, 1036-1040
 case reports, 1037-1040
 clinical characteristics of, 1036-1037, 1037-1040
 incidence of recurrences in, 1036-1040
 location of tumors in, 1036-1040
 frequency of, 1036
 histological findings in, 1037-1040
 results of treatment for, 1037-1040
 breast, duct epithelium, and fibroadenoma, method of labeling tissues with tritiated thymidine in vitro and its use in comparing rates of cell proliferation in, 639-643

Carcinoma(s), breast—(Contd.)

- female, metastatic, hormonal therapy with 26-methyl dihydrotestosterone propionate in, 705-706
- gastroduodenal metastases from, adrenal steroid-induced, report of, 426-431
- gelatinous, prognostic significance of, 699-704
- isocitric dehydrogenase, malic dehydrogenase, and glycolytic enzyme levels in patients with, 726-733
- metastatic, comparative evaluation of adrenalectomy and cortisone in treatment of, 1259-1265
- "recurrent" or inoperable, effect of hormonal treatment of, and patient survival in, report of, 300-307
- treated by 5-fluorouracil (5-FU), clinical evaluation of, 137-138, 140-145, 146-147, 148
- cervical, intraepithelial, microscopic study of margins of cone biopsy from, and predictability of residual carcinoma in uterus in, 795-800
- recurrent, after simple total hysterectomy, management of, 582-586
- dh59, use in experimental study of effectiveness of small amounts of preoperative irradiation in preventing growth of tumor cells disseminated at surgery, 284-295
- epidermoid, of larynx, incidence of cervical lymph node metastases from, and their relationship to certain characteristics of primary tumor, report of, 55-66
- esophageal, squamous cell, stromal inflammatory cell infiltration as prognostic factor in, 921-933
- gallbladder, dog, report of, 1127-1130
- use of methylcholanthrene as test carcinogen in, 1127-1130
- giant cell, of lungs, epithelial origin of confirmed by tissue culture, 1052-1056
- intraepithelial, uterine cervix, relative maturation of surface epithelium in, 1045-1051
- invasive, experimental, development of, from progression of cervical dysplasia in mouse, 201-204
- lung, declining rate of increase in mortality attributed to, 622-628
- treated with benzyl-N-bis(ethylene-imido)phosphorocarbamate (AB-103), report of, 853-861
- use of pancreatic deoxyribonuclease (dornase) in cytological diagnosis of, 419, 420
- oral, multiple, report on, 512-516
- cervical node involvement in, 514-515
- clinical features of abnormal oral epithelium in, 512-514
- atrophic areas in, 513
- clinical appearance of mucous membrane near primary lesions in, 513
- extrinsic factors in, 513
- intrinsic factors in, 513
- leukoplakia in, 513-514
- precancerous mucosal changes in, 513
- development of primary tumors at intervals in, 514
- of simultaneous primary tumors in, 514
- diagnostic procedures to be used in, 515-516
- division of tongue into geographical areas in, 516
- histopathology of, 512
- types of abnormalities found in, 512
- partial or hemi jaw resection in, 515
- patients with squamous cell carcinomas in, 512-516
- problems of correlation in, 515-516
- prosthetic rehabilitation in, 515
- radical neck dissection for involved cervical nodes in, 515
- salvage rates in, 515
- sites of primary tumors in, 514
- surgery or radiation therapy of primary tumor in, 515
- pancreatic, and multiple thrombi,

Carcinoma(s), pancreatic—(Contd.)

- morphological study of, 944-952
- small cell, cytological use of "wet films" in detection of, 1142
- squamous cell, as type of cancer of lung, validity and value of histological and cytological classifications in relation to, 673-698
- combined with trabecular adenocarcinoma, developing in dermoid cysts of ovary, report of, 990, 995-999
- in dermoid cysts of ovary, report of, 989-1000
- infiltrative, of vulva, 330-343
- intraepithelial, of vulva, 318-329
- multiple, oral, report of, 512-516
- of esophagus, stromal inflammatory cell infiltration as prognostic factor in, 921-933
- of lung, cytological use of "wet films" in detection of, 1140
- thyroid, in childhood, 734-743
- varying incidence of, in surgically removed thyroid glands, explanation of, 67-72
- undifferentiated, developing in dermoid cysts of ovary, report of, 990, 995-999
- large cell, as type of cancer of lung, validity and value of histological and cytological classifications in, 673-698
- of lung, cytological use of "wet films" in detection of, 1141
- small cell, as type of cancer of lung, validity and value of histological and cytological classifications in relation to, 673-698
- urokinase excretion in patients with, 889-896
- vulvar, infiltrative squamous cell (epidermoid), report of, 330-343
- intraepithelial and infiltrative, Bowen's type, report of, 318-329
- Carcinomatosis(es), lymphangitic, of lung, and high levels of amylase in body fluids, association of, 1109-1114
- pelvic and abdominal, primary in ovaries, attempted palliation by radical surgery in, 384-388
- Carcinosarcoma(s), cells, 256 Walker, from, in blood, effect of human fibrinolysin on survival of, 957-962
- in experimental studies of effect of nutrition on hepatic metastases, use of, 547-554
- in rats, influence of thermal stress and changes in body temperature on development of, 111-116
- CARR, A. H., *jr.*, *aw.*, 673-698
- CARTER, ROBERT E., and PIERCE, MILA I.: Studies on the relationship of viral infections to leukemia in mice, III. Failure to demonstrate acceleration of leukemia in mature AKR mice by cell-free brain filtrates from leukemic animals of the same strain, 496-501
- Catheterization, in antimetabolite-metabolite cancer chemotherapy using continuous intra-arterial 4-amino-N¹⁰-methyl pteroylglutamic acid (Methotrexate) with intermittent intramuscular citrovorum factor (CF), method of 744-752
- CAVANAGH, DENIS, *jr.*, *aw.*, 795-800
- Cell(s). *See also* Cytology
- and nucleus, diameters of, in study of measurement of quantitative characteristics of exfoliated cells from female genital tract, report of, 437-453
- argentaffin, in adenomatoid bronchial tumors, presence of, 171-173
- argyrophilic in adenomatoid bronchial tumors, presence of, 171-173
- atyphia of, from experimental cervical dysplasia in mouse, development of invasive carcinoma from, 201-204
- cancer, in operative wounds, significance of, 1041-1044
- association of lymph node metastases with local recurrence and distant metastases in, 1043
- data for local recurrences in, 1042-1043
- association of distant metastases with, 1042, 1043
- distant metastases, survival data, and, 1042
- time of local recurrence in, 1042

Cell(s), cancer—(Contd.)

- effect of lymph node metastases on wound-washing results in, 1042
- life table analysis in, 1041, 1042
- method of statistical analysis used in, 1041
- operative procedures among patients studied in, 1041
- results of study of, 1041-1043
- by anatomical sites of tumor in, 1041, 1042
- survival rates in, 1043
- carcinoma, embryonal, from teratocarcinoma, in vitro and in vivo study of differentiation in, 1017-1029
- carcinosarcoma 256 Walker, in blood, effect of human fibrinolysin on survival of, 957-962
- adult female white Wistar rats used in study of, 957
- aorta study after femoral vein infusion in, 960
- results at different time intervals in, 960
- fibrinolytic activity in, 959
- histological findings in, 961
- in vitro effect of fibrinolysin in, 958-959
- portal vein study after femoral vein infusion in, 960-961
- results at different time intervals in, 960-961
- preparation of tumor cell suspension in, 957-958
- testing of activity of human fibrinolysin in, 958
- by subcutaneous inoculations in, 958
- significance of findings in, 961-962
- vena cava study after femoral vein infusion in, 959
- results at different time intervals in, 959
- after jugular vein infusion in, 959-960
- results at different time intervals in, 960
- in rats, influence of thermal stress and changes in body temperature on, 111-116
- adrenocortical response in, 114, 115, 116
- appearance of tumors in, 113-114, 115
- application of "cold stress" in, 111-112, 113-114
- of "heat stress" in, 112, 114
- of hyperthermia in, 112-113, 114
- of hypothermia in, 112, 114
- definition of "cold stress" in, 111
- experimental method used in, 111
- results of, 113-115
- subcutaneous inoculation of tumor cells in, 113
- total plasma 17-hydroxycorticoid levels in, 114, 115, 116
- use of female Holtzman rats in, 111
- inoculation of rats with, effect of "aging" in reducing development of tumors after, 1057-1059
- cervical, cancer, exfoliated, interferometric study of anhydrous mass of, 378-383
- collection of, from female genital tract, methods of, as grouping factor in study of quantitative characteristics of exfoliated cells, 441
- epithelial, neoplastic, cytochemical study of esterase activity in, 369-377
- female genital tract, exfoliated, measurement methods and results of study of quantitative characteristics of, 437-454
- suitability of quantitative cytological measurements for automatic precreening of, 455-468
- from ganglioma of adrenal medulla, morphologically identical to hilus cells (extraparenchymal Leydig cells), report of, 421-425
- gastric, cancer, use of simplified chymotrypsin technique for exfoliative cytology in, 1074-1080
- giant, type of fibrosarcoma of brain, report of, 841-852

Cell(s)—(Contd.)

- hilus, similarity of, to cells from ganglioneuroma of adrenal medulla, report of, 421-425
- Kulschitzky, in adenomatoid bronchial tumors, distribution of, 172
- Leydig, extraparenchymal, similarity of morphologically to cells from ganglioneuroma of adrenal medulla, report of, 421-425
- malignant and normal, quantitative assessment of alkaline phosphatase activity in growth and differentiation patterns of, 502-506
- comparison of measurements in, 502-506
- for cytoplasm, 504, 505
- for heterochromatins, 504, 505
- for nucleoli, 504, 505
- equations used in, 504-505
- incubation in Gomori's substrate in, 502
- measurements of phase differences in, 502
- selection of cells in, 502
- staining techniques used in, 502
- use of Baker interference microscope in, 502
- mesenchymal, primitive, in histogenesis of malignant lymphomas, in African children, role of, 281-282
- normal and malignant, quantitative assessment of alkaline phosphatase activity in growth and differentiation patterns of, 502-506
- of spermatocytic seminoma of testis, structure of, 788-794
- oral, normal and malignant, exfoliated, effect of radiation on, 587-596
- lack of correlation between radiation response (AR) and clinical response of primary tumor in, 588-596
- between radiation response (AR) and radiation dose delivered in, 588-596
- between specific cytological changes and clinical response of primary tumor in, 588-596
- and radiation dose, 588-596
- palpable regional lymph nodes in, 590
- patients treated with ionizing radiation included in study of, 588
- presence of cancer cells in exfoliated material in, 589-592
- disagreement between clinical and cytological findings in, 591
- radiation response (AR) of normal cells in, 592-595
- results in, 588-596
- smears from surface of tumor and from adjacent normal mucosa in, 588
- criteria used to evaluate cells in, 588
- survival rates in, 589, 590
- tumor sites in, 590
- type of radiation therapy in, 588, 590
- proliferation of, comparison of rates of in tissues from duct epithelium, fibroadenoma, and carcinoma of breast, use of tritiated thymidine in, 639-643
- stromal inflammatory, infiltration by, in squamous cell carcinoma of esophagus, as prognostic factor, role of, 921-933
- trophoblastic, single, in routine vaginal smears, source of error in interpretation of, 1183-1185
- tumor, circulating, time of metastasis and release of, determined in experimental study, 883-888
- balance of factors between growing tumor and host reaction in, 886-887
- method of injection into mice in, 883-884
- possible explanation for presence of cells in mitotic division in peripheral blood in, 887
- sarcoma T241 Lewis used in, 883
- setup and results for experiment 1 (serial amputation of extremity bearing sarcoma T241 Lewis) in, 884, 885, 886-887

Cell(s), tumor—(Contd.)

- for experiment 2 (serial transfer of pulmonary tissue from mice bearing sarcoma T241 Lewis to peritoneal cavity of normal mice) in, 885, 886-887
- for experiment 3 (examination of venous blood from inoculated mice) in, 885, 886-887
- 3 methods of determining time of metastasis in, 883
- disseminated at surgery, effectiveness of small amounts of preoperative irradiation in preventing growth of, experimental study of, 284-295
- mesenchymal, cytochemical study of, esterase activity in, 369-377
- Central nervous system. *See also specific names of organs*
- gliomas of, as multiple primary malignant tumors of multicentric origin, report of, 244
- malignant lymphomas of, in African children, pathological entity of, 273-276
- Cervix. *See also Uterus*
- biopsies of including serial sectioning, critical study of, 213-216
- cases selected for restudy in, 214
- cervical specimens used in, 213
- cold knife conization material in, 214
- hysterectomy material in, 214
- multiple point biopsy material in, 214
- importance of carefully performed biopsy in, 215, 216
- possible reasons for disparity between cervical cell studies and subsequent biopsy diagnosis in, 213
- practicality of routine step sectioning for abnormal cases in, 213, 215-216
- preparation of specimens in, 214
- results of serial step sectioning in, 214-215
- cancer of, biometric approach to etiology of, 153-160
- classification into groups in, 154
- combinations of groups studied in, 157
- definition of variables under study in, 156
- distribution of clinic registrants by variables and by groups in, 154, 155
- of female clinic population by religion, age, marital events, and cancer status in, 156
- notation system used in separation of groups in, 156, 157
- population studied in, 154
- regression analysis in, 155-158
- simple correlations for each variable and multiple correlations for selected combinations of variables, with indications of most important variables, in study of, 157, 158-159
- summary of findings in, 158-160
- marital events as most important finding in, 158, 159-160
- use of frequency tables in, 155, 156, 157, 158-159
- variables studied in, 154
- experimental, interferometric study of anhydrous mass of exfoliated cells from, 378-383
- appearance of first recognizable cellular changes in, 380
- cell area and number of 20-methylcholanthrene paintings in fresh cells and alcohol-ether-fixed cells in, 382
- cellular vacuolation in dysplastic lesions in, 380
- collection of vaginal tumor cells in, 379
- comparison of fixed and unfixed normal cells in, 378
- cytoplasmic differences found in, 380-383
- induction of tumors by 20-methylcholanthrene in, 379
- nuclear area and number of 20-methylcholanthrene paintings in fresh cells and alcohol-

Cervix, cancer of—(Contd.)

- ether-fixed cells in, 382
- optical path difference in, 378
- percentage distribution of anhydrous nuclear mass in normal and abnormal cells in, 380
- of nuclear areas in normal and abnormal cells in, 379
- possibility of imbibition of water in, 381
- of role of mitochondrial enzymes in, 381-382
- preparation of smear specimens in, 379
- reading of measurements in, 379
- refractive index of medium in, 378
- of specimen in, 378
- relationship of anhydrous mass to number of 20-methylcholanthrene applications in, 381
- of cell area to nuclear area in basal cell hyperplasia in, 380
- in cancer cells in, 381-383
- in dysplasia in, 380-381
- of dry weight of object being measured in, 378
- of nuclear area to nuclear mass in normal and abnormal cells in, 379
- specific refractive increment in, 378
- thickness of object being measured in, 378
- type of Baker AO interference microscope used in, 379
- use of C3H female mice in, 379
- of interference microscope in, 378
- vaginal aspirations in late invasive lesions in, 380
- radiotherapy for, development of in situ epidermoid carcinoma of cervix and vagina after, 353-360
- case reports, 353-359
- changes in epithelial cells in, 359
- discovery of by routine cervical and vaginal smears in, 359, 360
- by routine random biopsy in, 359, 360
- histological appearance of recurrent in situ carcinoma in, 354-356, 358, 359
- initial biopsy diagnosis in, 353-359, 360
- operative procedures used in, 353-359
- postirradiation biopsy diagnosis in, 353-359, 360
- progression to invasive carcinoma in, 360
- radiation changes in cervicovaginal stroma and epithelium in, 359
- site of recurrence in, 353-359, 360
- time lapse between initial lesion and diagnosis of recurrence in, 353-359, 360
- type of radiation therapy used in, 353-359, 360
- carcinoma of, as multiple primary malignant tumors of multicentric origin, report of, 242
- recurrent, after simple total hysterectomy, management of, 582-586
- evaluation of radiation in, 585
- of surgical attack in, 585
- extensiveness of lesions in, 583
- interval between hysterectomy and first examination for recurrence in, 583
- prophylactic surgery less than 2 months after total hysterectomy in, 583, 585
- results according to pathological extension of tumor in, 583-585
- survival rates in, 583-585
- type of surgical procedures used in, 583-584
- of treatment given in, 582-584
- cells exfoliated from measurement method used in study of quantitative characteristics of, 437-453

Cervix—(Contd.)

- human, stratified squamous epithelium of, normal and cancer cells from, quantitative assessment of alkaline phosphatase, 502-506
- intraepithelial carcinoma of, microscopic study of margins of cone biopsy from and predictability of residual carcinoma in uterus in, 795-800
- cone margin prediction correlated with residual dysplasia found in hysterectomy specimen in, 797
- with residual intraepithelial carcinoma in hysterectomy specimen in, 797
- confirmation of diagnosis in, 795-796
- diagnosis based on pathological findings for hysterectomy specimen in, 796
- discrepancy between predicted and actual incidence of residual carcinoma in, 797-799
- division of patient groups in, 796
- method of evaluating slides in, 795-796
- technique of preparing cone biopsy specimen in, 796
- transection of deep endocervical glands correlated with residual carcinoma in hysterectomy specimen in, 797
- lymph node metastases to, from epidermoid carcinoma of larynx, incidence of and relationship to certain characteristics of primary tumor, report of, 55-66
- mouse, progression of experimental dysplasia in, 201-204
- development of dysplasia after, 201-202
- types of changes seen in cells histologically in, 201-204
- of invasive carcinoma after, 202-204
- preparation of cell studies in, 201
- use of female C3H mice in, 201
- application of methylcholanthrene in, 201
- suitability of quantitative cytological measurements for automatic pre-screening of cells exfoliated from, report of, 455-468
- uterine, human, cancer of, studies on heredity factor in, 179-194
- accuracy of patient diagnosis in, 181
- age of patients and controls in, 183, 185
- comparison of cervical cancer patients, controls, and general California population in, 180
- completeness of pedigrees in, 181-183, 184
- estimate of completeness by use of "twin set loss" test in, 182, 183
- data for cancer rates in twin sets in, 186-187, 193
- on pedigree studies in, 187-194
- environment versus heredity in, 187, 189, 189-192
- distribution of cancer by site and sex in families of cancer patients and controls in, 186, 192, 193, 194
- follow-up in, 180-181
- code used in, 181
- use of "statistical cancer pedigree" in, 180, 181
- geographical origin of samples in, 181
- incidence of cancer in families of propositi in, 183-193
- comparison of total sisters and of sisters to brothers in, 185, 191
- frequencies in male and female relatives in, 185-189
- rates by tumors with epidermoid cellular structure in, 183, 186, 193
- for single and multiple case families in, 184-185, 188-190, 193-194
- for sisters of controls in, 190
- of skin cancer among, 186-187, 192, 193

Cervix, uterine—(Contd.)

- patient sample used in, 179-180
- race of patients and controls in, 181
- religion of patients and controls in, 181, 182
- selection of controls in, 179-180
- age as factor in, 179-180
- race as factor in, 180
- religion as factor in, 180
- size of samples and families in, 181, 182
- type of questionnaire used in, 180
- uterine and cervical cancer incidence in sisters of cancer patients and those of controls in, 186, 193
- metaplasia, dysplasia, and intraepithelial carcinoma of, relative maturation of surface epithelium in, 1045-1051
- criteria used in, 1046
- hematoxylin and eosin-stained sections used in, 1045-1046
- percentage of each sample area occupied by nuclei in, 1047-1051
- comparisons of, in various layers of epithelium, in, 1047-1050
- selection of samples of tissue for analysis in, 1046-1047
- spread of cancer in, as seen in giant histological sections, report of, 42-54
- extension to level above internal cervical os in, 46, 53
- ganglion cells in, 46, 52
- identification of lymph nodes in, 50
- method of processing tissues in, 42-45
- mounting of sections in, 42-43
- patterns found in cases of direct extension into vagina in, 50-51
- previous irradiation and, 50
- spread into fundus in, 51
- types of histological features tabulated in, 45
- of tumor extension in, 45-54
- in anterior exenteration specimens, 45
- in paracervical tissue, 45-46
- in paravaginal tissue, 46
- in perineural tumor, 46
- in Wertheim hysterectomy specimens, 45
- ureteral involvement in, 46-50
- Cv. See *Citrovorum* factor
- CHAN, P. Y. M.; HADDEN, D. H.; McDONALD, G. O., and COLE, W. H.: The use of "aging" in reducing the development of tumors after inoculation of carcinosarcoma 256 Walker cells, 1057-1059
- CHATTERJEE, R., *jt. aw.*, 502-506
- CHAUDHRY, ANAND P., *jt. aw.*, 73-101
- Chemical(s). *See also specific names*
- carcinogenic, in newborn mice, effects of, 308-317
- association of malignant lymphomas with pulmonary tumors in, 315
- histiocytic ("reticulum cell") type of, 314
- liver involvement in, 313
- medium-sized lymphocyte (lymphocytic) type of, 314
- undifferentiated reticular cell (stem cell) type of, 313-314
- control groups used in, 308
- injection of aqueous gelatin in, 308
- injection of noncarcinogenic polycyclic hydrocarbon 9-methylanthracene (MeA) in, 308
- incidence of malignant lymphoma in, 308-316
- average latent period before appearance of, 309, 310
- of pulmonary tumors in, 308
- injection of compounds in, 308
- miscellaneous neoplastic lesions in, 315
- pathological findings for malignant lymphoma in, 310-315
- histological criteria for diagnosis in, 310-313

Chemical(s), carcinogenic—(Contd.)

- pattern of infiltration in lungs in, 313
- polycyclic hydrocarbons in, 308
- benzo[a]pyrene (BP) as, 308
- dibenzo[a,h]anthracene (DBA) as, 308
- 7,12-dimethylbenzo[a]anthracene (DMBA) as, 308
- 3-methylcholanthrene (MC) as, 308
- study of ethyl carbamate (urethan) in, 308
- use of Swiss albino mice in, 308
- effect of, on radiation sensitivity of mouse tumors, 117-121, 124
- Chemotherapy. *See also specific names of agents*
- 6-aminonicotinamide (6-AN), in therapy of disseminated cancer, clinical use of, 644-648
- chlorambucil in treatment of childhood cancer, use of, 36-41
- combined with surgery, in treatment of malignant tumors, results of, 1197-1204
- effect of intrathecal cyclophosphamide (Cytosan) in dogs, 1115-1116
- experimental, using Methotrexate (amethopterin, 4-amino-N¹⁰-methyl pteroylglutamic acid), growth rates of transplanted human cancer during, 1117-1121
- for giant follicular lymphoblastoma, report of, 771, 772-773
- in cancer, with 5-fluorouracil (5-FU), clinical evaluation of, 131-152
- of cancer, distribution of naphthylchloracetamide and, 296-299
- prolonged intra-arterial technique for inoperable cancer, 493-495
- use of continuous intra-arterial 4-amino-N¹⁰-methyl pteroylglutamic acid (Methotrexate) with intermittent intramuscular citrovorum factor (CV), report of, 744-752
- Chlorambucil, in therapy of giant follicular lymphoblastoma, report of, 771, 772-773
- use in childhood cancer, of, 36-41
- chemical structure of, 36
- Choline, levels of, in experimental studies of effect of nutrition on hepatic metastases, report of, 550-553
- Cholinesterase(s), reaction, in juvenile melanoma, report of, 1002, 1003
- Chondrosarcoma(s), in children, as part of malignant compound tumor, report of, 524-533
- CHRISTOPHERSON, WILLIAM M., and BROGHAME, WALTER, JR.: Progression of experimental cervical dysplasia in the mouse, 201-204
- and PARKER, JAMES E.: A critical study of cervical biopsies including serial sectioning, 213-216
- jt. aw.*, 378-383
- Chromotrope 2R, staining of sections from rats with nutritional cirrhosis with, 963-970
- CHU, ELIZABETH W., *jt. aw.*, 883-888
- CHU, FLORENCE C. H., and HILARIS, BASIL E.: Value of radiation therapy in the management of intracranial metastases, 577-581
- Chymotrypsin, in exfoliative cytology of stomach cancer, report of, 1074-1080
- Cigarette(s). *See also Smoking and Tobacco*
- microscopic properties of basement membrane and elastic fibers of trachea and bronchus in relation to, 1175-1182
- role of acidic fractions of, as promoters in carcinogenesis, study of, 1306-1315
- tar, and beta radiation, in mice, combined action of, 344-352
- appearance of erythema and epilation in areas of radiation exposure in, 345
- of leukemia in, 346, 347-348
- of lung tumors in, 346-347
- alveolar carcinoma type of, 346
- of mammary gland tumors in, 347
- of metastatic lesions of lung tumors and mammary gland tumors in, 347, 348, 350
- of ovarian tumors in, 348-349, 351

- of skin cancer in, 345-346
basic fraction obtained in, 344
causes of death of mice in, 350
comparison of effect of beta radiation alone in, 349, 350-352
of tar alone in, 349, 350-352
development of hepatomas in, 348, 351
dosage of beta radiation used in, 344
groups of mice used in, 344-345
tar painting and radiation exposure plans used in, 345
incidence of multiple tumors in, 348, 350
life spans of animals in different groups in, 350, 352
method of administering beta radiation in, 344
microscopic appearance of mouse organs in, 349
other types of cancers appearing in, 348-349
painting of mice in, 344
smoking machine used in, 344
strontium 90, RA-2A, medical applicator used for administering beta radiation in, 344
Cirrhosis. *See also* Hepatectomy, Liver, and Metastasis(es)
activity of triphosphate isomerase in blood serum from patients with, 1166-1174
nutritional repair of, autoradiographic and histological study after partial hepatectomy in, 963-970
autoradiography and mitotic counts at different time intervals in, 967-969
estimation of liver regeneration in, 965-970
occurrence of, throughout parenchyma, in, 965-970
definition of cirrhosis as, 969
use of different stains to demonstrate fibrous tissue as, 969
variation in cirrhosis in different animals as, 969
in cirrhosis in same lobe of liver as, 969
fixation of liver sections in, 965
stains used in, 965
chromotrope 2R as, 965
labeling of cells with tritium (H^3) in, 965
histological findings in, 965-966
grading of degree of cirrhosis in, 965
variation in degree of cirrhosis in, 964, 965
method of determining percentage of partial hepatectomy in, 965
use of Sprague-Dawley male rats in, 963
Citrate, infusion of, in advanced cancer patients, effect of on renal phosphate and calcium excretion, 934-943
Citrovorum factor (CF), intermittent intramuscular, and 4-amino- N^{10} -methyl pteroylglutamic acid (Methotrexate), in antimetabolite-metabolite cancer chemotherapy, report of, 744-752
CLARK, KEMP, *jt. au.*, 1115-1116
CLARKSON, BAYARD D., *jt. au.*, 744-752
Classification, cytological, of cells exfoliated from female genital tract, quantitative characteristics of, 441-442
CLIFFORD, PETER, *jt. au.*, 744-752
CLIFFORD, EUGENE E., and GROSSI, CARLO E.: Pancreatic deoxyribonuclease (dornase) aerosol in treatment of bronchopulmonary complications and tracheitis sicca, 414-420
jt. au., 957-962
COHEN, RICHARD B., *jt. au.*, 421-425
Cold, as physical agent affecting radiation sensitivity of mouse tumors, report of, 122-123
COLLE, WARREN H., *jt. au.*, 111-116, 1057-1059
Colon. *See also* Anus and Rectum
adenocarcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 240
primary cancer of, frequency of occurrence of second primary malignant tumors in women with, 232
CONNAUGHTON, PETER J., and LEWIS, F. JOHN: The cancericidal effects of X irradiation at body temperature below $10^{\circ}C$, 1060-1062
COON, WILLIAM W., *jt. au.*, 1081-1084
CORNE, JOHN S.: Multiple lymphomatous polyposis of the gastrointestinal tract, 249-257
Cortisone, and adrenalectomy, therapy with, for advanced breast carcinoma, evaluation of, 1259-1265
in therapy of giant follicular lymphoblastoma, 771, 772-773
COTRAN, RAMZI S.: Metastasizing basal cell carcinomas, 1036-1040
COWDREY, E. V.; CRONINGER, A.; SOLARIC, S., and SUNTZEFF, V.: Combined action of cigarette tar and beta radiation on mice, 344-352
jt. au., 1175-1182
"CPL" classification, in evaluating prognosis in squamous cell carcinoma of esophagus, use of, 925-932
CREEP, ANDRE G., *jt. au.*, 1151-1165
CRONINGER, A., *jt. au.*, 344-352
Crystalloid(s), of Reinke, in ganglioneuroma of adrenal medulla containing cells morphologically identical to hilus cells (extraparenchymal Leydig cells), report of, 421-425
Cyclophosphamide (Cytosan), intrathecal, in dogs, effect of, 1115-1116
dose injected in, 1115-1116
method of anesthetizing in, 1115
pH of drug in, 1115
purpose of study of, 1115
results in, 1116
by amounts of dose in, 1116
neurological symptoms in, 1116
symptoms of toxicity in, 1116
(Endoxan), in study of combined surgery and chemotherapy, use of, 1197-1204
Cylindroma(s), bronchial, nonbenign nature of, 539-546
Cyst(s), dermoid, of ovary, cancer developing in, 989-1000
case reports, 900-994
structure in relation to types of cancer that may develop in, 989
Cysteine, effect of, on radiation sensitivity of mouse tumors, 118, 119, 120
Cyt analyzer, in study of suitability of quantitative cytological measurements for automatic prescreening of cells exfoliated from female genital tract, report of, 467
Cytology. *See also* Cell(s)
exfoliative, of renal pelvic lesions, use of, 1085-1087
measurement methods in study of quantitative characteristics of exfoliated cells from female genital tract, report of, 437-453
suitability of quantitative cytological measurements for automatic prescreening of exfoliated cells from female genital tract, 455-468
vaginal, histiocytes and histiocytic reaction in, survey of risks of false positive diagnosis in, 1223-1225
Cytosan (cyclophosphamide), intrathecal, in dogs, effect of, 1115-1116
DANIEL, WILLIAM W., and BRUNSCHWIG, ALEXANDER: The management of recurrent carcinoma of the cervix following simple total hysterectomy, 582-586
jt. au., 353-360
DAO, THOMAS L.; TAN, EDWARD, and BROOKS, VIVIAN: A comparative evaluation of adrenalectomy and cortisone in the treatment of advanced mammary carcinoma, 1259-1265
DBA (dibenz[*a,h*]anthracene), as carcinogenic chemical, in newborn mice, report of, 308-317
DE, P.; CHATTERJEE, R.; BHATTACHARYA, K., and MITRA, S.: Quantitative assessment of alkaline phosphatase activity in growth and differentiation patterns of normal and malignant cells, 502-506
Dehydrogenase(s), isocitric (icd), levels of, in serum of patients with carcinoma of breast, 726-733
malic (md), levels of, in serum of patients with carcinoma of breast, report of, 726-733
DENBOW, VICTOR D., and ADAIR, FRANK E.: Lymphangiosarcoma in the postmastectomy lymphedematous arm. A case report of a 10-year survivor treated by interscapulothoracic amputation and excision of local recurrence, 210-212
DENK, W., and KARRER, K.: Combined surgery and chemotherapy in the treatment of malignant tumors, 1197-1204
Dentinoma(s), in man, classification, histopathology, and clinical behavior of, 86-87
Deoxyribonuclease, pancreatic (dornase), aerosol, in treatment of bronchopulmonary complications and tracheitis sicca, report of, 414-420
case reports, 416-417
complications after, 419
indications for use of, 415
previous experience with, 414
results in chronic bronchopulmonary disease in, 418-419
in cytological diagnosis of carcinoma of lung in, 419
in postoperative atelectasis and postoperative pulmonary complications in, 415-416
in tracheitis sicca in, 418
use prior to bronchoscopy, 419
Deoxyribonucleic acid (DNA), increase in, in epithelial cells of mouse kidney infected with SE polyoma virus, study of, 570, 573
Dermatitis, atopic, complicated by reticular cell sarcoma, prolonged remission after Potaba therapy, report of, 5-12
Dibenz[*a,h*]anthracene (DBA), as carcinogenic chemical, in newborn mice, report of, 308-317
2,2-Dichloro-*N*-methyl-*N*-oxide hydrochloride (Mitomen, Nitromin), in combined surgery and chemotherapy as treatment for malignant tumors, use of, 1197-1204
Diet, as etiological factor in cancer of esophagus, study of, 389-413
choline-deficient, as cause of cirrhosis in rats, autoradiographic and histological study of repair after partial hepatectomy, 963-967
effect of, on hepatic metastases, experimental studies of, 547-554
Differentiation, of teratocarcinomas, *in vitro* and *in vivo* study of, 1017-1029
patterns of, in normal and malignant cells, quantitative assessment of alkaline activity in, 502-506
Digestive tract. *See specific names of organs*
1,4-Dimethanesulfonoxymethane, radioactive-labeled (C^{14} -labeled Myleran or busulfan and S^{35} -labeled Myleran), excretion and distribution studies in cancer patients with, 953-956
7,12-Dimethylbenz[*a*]anthracene (DMBA), as carcinogenic chemical, in newborn mice, report of, 308-317
role of in carcinogenesis, experimental study of, 1306-1315
Dissection, neck, radical, and primary en bloc laryngectomy, for epidermoid carcinoma of larynx, incidence of cervical lymph node metastases found in, 55-66
DIXON, W. J., *jt. au.*, 152-160
DMBA (7,12-dimethylbenz[*a*]anthracene), as carcinogenic chemical, in newborn mice, report of, 308-317
role of in carcinogenesis, experimental study of, 1306-1315
DNA. *See* Deoxyribonucleic acid
DOCKERTY, MALCOLM B., *jt. au.*, 221-230, 231-237, 238-248, 901-912
Dog(s), in study of effect of intrathecal cyclophosphamide (Cytosan), report of, 1115-1116
mongrel, development of carcinoma of gallbladder in, 1127-1130
small intestine of, effect of succinyl-sulfathiazole (Sulfasuxidim) on radiation response of, 862-874
Dornase (pancreatic deoxyribonuclease), aerosol, in treatment of bronchopulmonary complications and tracheitis sicca, report of, 414-420
Dornase (pancreatic deoxyribonuclease) in treatment of bronchopulmonary complications and tracheitis sicca, report of, 414-420
DROLETTE, HERNADETTE M., *jt. au.*, 330-343

- Drug(s), alkylating, radioisotope-labeled, in cancer patients, excretion and distribution studies of, 953-956
dosage for each compound used in, 954
location of isotopic element employed in, 953
molecular structure of compounds used in, 953
procedure followed in, 954
radioactivity measurement methods in, 954
urinary collections in, 954
results of individual experiments in, 954-955
circulating radioactivity levels in blood in, 955
specific activities of lymph node and muscle at biopsy after, 955, 956
time factor in, 955-956
urinary excretion of radioactivity levels in, 955
use of 1,4-dimethanesulfonylbutane (C^{14} -labeled Myleran or busulfan and S^{35} -labeled Myleran) in, 953-956
of triethylene melamine (C^{14} -labeled TEM) in, 953-956
of triethylenephosphoramide (P^{32} -labeled TEPA) in, 953-956
"Dual antagonist," benzyl-N-bis(ethylenimido)phosphor-carbamate (AB-103) as, in treatment of bronchogenic carcinoma, use of, 853-861
Dualar. *See* Benzyl-N-bis(ethylenimido)phosphor-carbamate
DUDLEY, H. ROBERT, *jr.* *see*, 517-523
DUFF, JOHN K.; SULLIVAN, ROBERT D.; MILLER, EDWARD; ULM, AARON H.; CLARKSON, BAYARD D., and CLIFFORD, PETER: Antimetabolite-metabolite cancer chemotherapy using continuous intra-arterial Methotrexate with intermittent intramuscular citrovorum factor. Method of therapy, 744-752
Duodenum, metastases to, from carcinoma of breast, adrenal steroid-induced, report of, 426-431
DUTZ, WALTER, and STOUT, ARTHUR PURDY: The myxoma in childhood, 629-635
Dye(s), naphthol AS acetate as, in cytological study of esterase activity in human neoplasms and stromal macrophages, report of, 369-377
radio-opaque, injection of, in measurement studies of tumor growth by lymphangiography of retroperitoneal lymph nodes, 205-209
sky-blue, in anatomical studies of dissemination of cancer of larynx, report of, 1131-1138
Dysplasia(s), cervical, mouse, experimental, progression of, to invasive carcinoma, study of, 201-204
application of methylcholanthrene in, 201
cell changes seen histologically after, 201-204
development of invasive carcinoma from, 202-204
in bronchial mucosa, histological study of, 1226-1234
previous studies of, 1226, 1231-1234
relationship of age to, 1226-1234
of sex to, 1226-1234
of smoking to, 1226-1234
of uterine cervix, relative maturation of surface epithelium in, 1045-1051
E-39 (2,5-bis(ethylenimino)-3,6-dipropoxy-1,4-benzoquinone), in combined surgery and chemotherapy as treatment for malignant tumors, use of, 1197-1204
EDDY, BERNICE E., *jr.* *see*, 567-576
EFFLER, DONALD B., *jr.* *see*, 1139-1143
ENGE, NORMAN: Amylase activity in body fluids, 1109-1114
Endocrine(s). *See also specific names*
in luteinization of stroma in metastatic ovarian cancer, 827-840
Endoxan (cyclophosphamide) in combined surgery and chemotherapy, use of, 1197-1204
Environment. *See also* Epidemiology
as basis for clinical syndrome of malignant lymphoma in African children, possibility of, 267
Environment—(Contd.)
as etiological factor in cancer of esophagus, study of, 389-413
epidemiological study of seasonal incidence of acute leukemia, 1301-1305
incidence of cancer in Singapore children, report of, 534-537
of malignant lymphoma in African children, role of, 270-283
influence of thermal stress and body temperature changes on development of carcinosarcoma 256 Walker cells in rats, study of, 111-116
possible role of, in incidence of stomach cancer in spouses of patients with, report of, 199
versus heredity, in cancer of human uterine cervix, studies on, 187, 189, 189-192
Enzyme(s). *See also specific names*
activity of, in human blood serum from normal individuals and from individuals with various disease conditions, 1166-1174
of naphthylchloroacetamide in mouse tissue and human organs and serum, 296-299
in juvenile melanoma, histochemical study of, 1001-1004
levels of, in serum of patients with breast carcinoma, 726-733
pancreatic deoxyribonuclease (dornase), as, in treatment of bronchopulmonary complications and tracheitis sicca, report of, 414-420
complications from, 419
indications for use of, 415
Epidemiology. *See also* Statistics and Environment
analysis of 5 "stomach cancer families" in State of Utah, 1005-1016
biometric approach to etiology of cancer of cervix, 153-160
clinical syndrome of malignant lymphomas in African children, report of, 258-269
declining rate of increase in mortality attributed to carcinoma of lung, report of, 622-628
etiological factors in cancer of esophagus, study of, 389-413
incidence of cancer in Singapore children, report of, 534-537
of stomach cancer in spouses of patients with, 199-200
pathological entity of malignant lymphoma in African children, report of, 270-283
role of acidic fractions of tobacco as promoters in carcinogenesis, study of, 1306-1315
seasonal incidence of acute leukemia, study of, 1301-1305
studies on inheritance of cancer of human uterine cervix, 179-194
Epilation, after exposure of mice to combined action of cigarette tar and beta radiation, appearance of, 345
Epithelium(s), as origin of giant cell carcinoma of lungs, confirmation of by tissue culture studies, report of, 1052-1056
duct, breast carcinoma, and fibroadenoma, method of labeling tissues with tritiated thymidine in vitro and its use in comparing rates of cell proliferation in, 639-643
of basement membranes of trachea and bronchus, in smokers and nonsmokers, results of measurements of, 1175-1182
oral, abnormal, in cases of multiple primary oral carcinomas, clinical features of, 512-514
surface, in metaplasia, dysplasia, and intraepithelial carcinoma of uterine cervix, relative maturation of, 1045-1051
tubular, of mouse kidney, infected with SE polyoma virus, and stroma, difference in host cell-virus relationship in, 567-576
Epulis, pigmented, of infancy, congenital melanocarcinoma, melanotic adaman-tinoma, retinal anlage tumor, and prog-noma, report of case in adult, 1090-1108
Erratum, 1150
Erythema, after exposure of mice to combined action of cigarette tar and beta radiation, appearance of, 345
Eserine sulfate, inhibitory effect of, on esterase activity of human neoplasms and stromal macrophages, 373, 375
Esophagus, cancer of, study of etiological factors in, 389-413
alcohol consumption in, 395-396
amount of drinking in, 395
analysis by religion for drinkers in, 396
standardization for smoking of drinkers in, 395-396
type of alcohol consumed in, 395
background data for American female patients in, 400-402, 403
age in, 400
alcohol consumption in, 401, 403
amount of drinking in, 401, 403
type of alcohol consumed in, 401, 403
dental factors in, 401
dietary factors in, 401
butter consumption in, 401
citrus fruit consumption in, 401
egg and green and yellow vegetable consumption in, 401
milk consumption in, 401
educational status in, 400
marital status in, 400
race in, 400
religion in, 400
smoking habits in, 400-401, 403
for American male patients in, 393-400, 401, 402
environmental habits of smokers and nonsmokers in, 394
histories for male nonsmokers in, 395
place of birth in, 394
race in, 393
religion in, 393
smoking habits in, 394-395
amount smoked in, 394
length of cigarette or cigar smoked in, 394
tobacco chewing in, 394
standardization for alcohol consumption of smokers in, 394
for religion of smokers in, 395
type of residence in, 393-394
changes in incidence in, 390
dental factors in, 396
dietary factors in, 396-397
egg and green and yellow vegetable consumption in, 396-397, 400
milk and butter consumption in, 396-397, 400
epidemiological pattern in, 408-410
data for Africa in, 409
for India in, 409-410
incidence of multiple primaries in, 410
low income status in, 410
other upper alimentary and respiratory cancers in, 410
for Western European countries in, 409
incidence rates in, 408-409
religion in, 409
sex ratio in, 409
geographical factors in, 389-390
death rates by, 389-390
histories for nondrinkers in, 398
methodology used in, 392-393
age distribution in, 393
educational status in, 393
heredity in, 392
income group in, 390
interview method used in, 392
marital status in, 393
patients and controls in, 392
sex differences in, 390
tylosis in, 392
miscellaneous factors in, 397-400
blood types and, 399
body build and, 398
cross analysis of milk and alcohol consumption among men in, 400
family history and, 399

Esophagus, cancer of—(Contd.)

- food temperature and, 398
- history of previous diseases in, 397
- location of lesion in, 397
- multiple primary cancers among men with esophageal cancer in, 401
- occupational data and, 397
- statistical summary table for American male group in, 402
- summation of data for Indian patients in, 403-405, 406
- alcohol consumption in, 404, 406
- background data in, 403
- dietary factors in, 404-405
- miscellaneous factors in, 404-405, 406
- statistical summary table for, 406
- tobacco usage in, 403-404
- bidi smokers in, 403-404
- quid chewing in, 404, 405
- type and amount of tobacco consumed in, 404
- for Swedish patients in, 402-403
- percentage distribution of Plummer-Vinson syndrome symptoms in, 402, 403
- tobacco and alcohol consumption in, 403
- suspected etiological factors in, 390-392
- air pollution as, 392
- alcohol as, 391
- cidiviki as, 391
- pai kae as, 391
- betel quid chewing as, 390
- diet as, 391
- correlation of Plummer-Vinson disease incidence and, 391
- incidence of double primaries as, 392
- kaffir pipe smoking as, 390
- ratio of observed to expected death rates for men in, 390-391
- oral hygiene as, 391
- stricture due to lye exposure as, 392
- syphilis as, 391
- thermal effects of hot food and drinks as, 391-392
- tobacco as, 390
- carcinoma of, squamous cell, stromal inflammatory cell infiltration as prognostic factor in, 921-933
- average invasive power and survival time in, 931
- cell type in, 922-924
- anaplastic, 923
- basal, 922-923
- noncornified squamous, 922
- well differentiated squamous, 922
- definition of groups in, 921
- group A (patients who recovered from and lived more than 3 months postoperatively) in, 921
- subdivision of group A in, 921
- group B (patients whose deaths were attributed directly to surgery or occurred within 2 months of operation) in, 921
- group C (patients who underwent surgery and upon whom autopsy was done in, 921
- evaluation of surgical treatment by, 929-932
- features of prognosis in, 924-927
- relationship of cell type of tumor and end results in, 924, 925
- "CPL" classification system used in, 925, 926
- of grade of tumor and end results in, 924, 925-926
- of lymph node metastases and end results in, 926-927
- of site of tumor and end results in, 923, 924
- of size of tumor and end results in, 924, 925
- intraepithelial carcinoma (carcinoma in situ) in, 923, 924
- macroscopic features in, 922
- diffuse infiltrating type of tumor in, 922

Esophagus, carcinoma of—(Contd.)

- length of tumors in, 922
- sessile polypoid structure type of tumor in, 922
- solitary or multiple tumors in, 922
- ulcerating infiltrating type of tumor in, 922
- microscopic features in, 922-924
- mode of spread in, 923-924
- destructive or "regressive" changes in tumor cells in lymph nodes in, 924
- numerical evaluation of invasive power in, 926-927, 928-929, 930-931
- review of clinical material in, 922-927
- age distribution in, 922
- peak incidence and, 922
- distribution of tumors by site in, 922
- duration of symptoms in, 922
- male and female ratio in, 922
- type of symptoms in, 922
- stromal reaction in, 927-929
- relationship of, to end results in, 925, 926
- biological aggressiveness of tumor in, 927
- definition of invasive power (IP) in, 927
- coefficient of invasive power and defensive power (DP) in, 928-929
- host defense in, 927-928
- evaluation system for, 927
- surgical material included in study of, 921
- epitheliomas of, as multiple primary malignant tumors of multicentric origin report of, 239
- Esterase(s), activity of, in human neoplasms and stromal macrophages, cytochemical study of, 369-377
- activity at cell boundaries in, 374, 375
- at luminal or basal aspects of cells in, 371, 374, 375
- in cell cytoplasm of, 369-377
- tumor stroma in, 372-373
- breast duct cells in, 373
- collagenous reticular fibers in, 373
- eosinophils, lymphocytes, and plasma cells in, 373
- fat cells in, 373
- fibroblasts in, 373
- ganglion cells in, 373
- giant cells in, 373
- histiocytes in, 372
- intestinal surface epithelium, goblet cell theca, and columnar cells in, 373
- liver parenchymatous and renal tubule cells in, 373
- macrophages in, 372-373
- mast cells in, 373
- Paneth's cells in, 373
- parietal and pyloric gland cells in stomach in, 373
- plasma in, 373
- polymorphonuclear leukocytes in, 373
- red cells in, 373
- smooth muscle fibers in, 373
- by type of benign neoplastic cells, 373
- of malignant neoplasm, 370-372
- comparison of activity with different substrates being used in, 370
- control reactions used in, 369
- identification of macrophages in, 375
- inhibitory effect of eserine sulfate in, 373, 375
- of sodium fluoride in, 373, 375
- of sodium taurocholate in, 373, 375
- preparation of tissues in, 369
- use of fast garnet salt GBC in, 369
- of naphthol AS acetate as substrate in, 369
- spheroidal bodies seen in, 375
- stains used in, 369
- tissue blocks used in, 369
- variations of intensity and appearance of color reaction in, 375
- Estrogen(s). *See also specific names* in treatment of primary malignant tumors, development of second primary malignant tumor after, 231-232, 236

- Ethyl carbamate (urethan), as carcinogenic chemical, in newborn mice, report of, 308-317
- Etiology, of cancer of esophagus, study of factors in, 389-413
- of pure myxoma in childhood, 631
- Excision, local, as treatment for adenoid cystic carcinoma, inadequacy of, 1247-1250
- radical, as treatment for adenoid cystic carcinoma, report of, 1248
- Exfoliation, of normal and malignant oral cells, effects of radiation on, 587-596
- Eye(s), involvement of, in giant follicular lymphoblastoma, report of, 757-758
- Fast garnet salt GBC, in cytochemical study of esterase activity in human neoplasms and stromal macrophages, report of, 369-377
- Fat, content of, in diet, effect of on hepatic metastases, experimental studies of, 547-554
- Female genital tract. *See specific names of organs*
- Fever, incidence of, in patients with giant follicular lymphoblastoma, report of, 763
- Fiber(s), elastic, and basement membrane, of trachea and bronchus, of smokers and nonsmokers, microscopic properties of, 1175-1182
- Fibrinolysin, human, effect of, on survival of carcinosarcoma 256 Walker cells in blood, report of, 957-962
- Fibroadenoma(s), breast carcinoma, and duct epithelium, method of labeling tissues with tritiated thymidine in vitro and its use in comparing rates of cell proliferation in, 639-643
- Fibroma(s), ameloblastic, in man, classification, histopathology, and clinical behavior of, 82-85
- cementifying, in man, classification, histopathology, and clinical behavior of, 90-95
- odontogenic, in man, classification, histopathology, and clinical behavior of, 89, 90
- Fibrosarcoma(s), giant cell, of brain, report of, 841-852
- FIDLER, H. K., *jr.* *et. al.*, 1226-1234
- FISHER, BERNARD, and FISHER, EDWIN R.: Experimental studies of factors influencing hepatic metastases. VI. Effect of nutrition, 547-554
- FISHER, EDWIN R., *et. al.*, 547-554
- Florisil, in chromatography of 7,12-dimethylbenz[a]anthracene, use of, 1307
- Fluid(s), body, amylase activity in, 1109-1114
- case reports, 1111, 1112
- clinical diagnosis of pneumococcal empyema and, 1109
- coincidence of alcoholism and, 1109, 1110, 1113
- of cirrhosis of liver and, 1109, 1110, 1113
- definition of negative, borderline, and positive cases in, 1109
- levels in cases of carcinoma of lung in, 1109-1114
- method of determination used in, 1109
- neoplasms associated with high levels in, 1109-1111
- Papanicolaou classifications in, 1110, 1111
- role of pancreatitis in, 1109-1114
- significance of high levels and lymphangitic carcinomatosis in, 1114
- site of fluid in, 1110-1112
- types of disease and high levels in, 1111
- Fluorescein (Fluorescite) as dye used in antimetabolite-metabolite cancer chemotherapy using continuous intra-arterial 4-amino-N¹⁰-methyl pteroylglutamic acid (Methotrexate) with intermittent intramuscular citrovorum factor (CF), report of, 747
- Fluorescite. *See Fluorescein*
- Follicle(s), hair, role of, in origin and evolution of benign epithelial tumors, report of, 801-816
- FOOTE, FRANK W., *Jr.*, *et. al.*, 699-704
- FORAKER, ALVAN G., *et. al.*, 1045-1051
- FORD, D. K.; FIDLER, H. K., and LOCK, D.R.: Dysplastic lesions of the bronchial tree, 1226-1234

- FORNER, CLAUDE E., JR., *jr. sm.*, 608-621
- FORTNER, JOSEPH G., and LEFFALL, LA-SALLE D.: Carcinoma of the gallbladder in dogs, 1127-1130
- FOURNIER, PIERRE, *jr. sm.*, 817-826
- FREI, EMIL, III, *jr. sm.*, 608-621
- FRICKELL, GILBERT H., and PARSONS, LANGDON: The spread of cancer of the uterine cervix as seen in giant histological sections, 42-54
- SHERMAN, JOSEPH D., and SOMMERS, SHELDON C.: Growth curves of human cancer transplants during experimental chemotherapy, 1117-1121
- 5-Fluorouracil (5-FU), cancer chemotherapy with, clinical evaluation of, 131-152
- case reports, 139, 146-151
- classification of toxicity in, 132-133
- clinical evaluation of regression in, 132
- comparison of total doses tolerated by patients with and without regressions who did not receive radiation in, 146
- complications of toxicity in, 133-135
- correlation between dosage, performance status, previous therapy, and improvement in, 137
- and toxicity in, 136-137
- duration of regressions in, 140-145
- factors aggravating toxicity in, 135-136
- incidence of toxicity in, 132, 133
- changes in myeloid-erythroid ratio compared with degree of leukopenia as, 133, 134
- manifestations of, 132, 133
- over-all therapeutic results achieved in, 136-151
- results for adenocarcinomas of large bowel in, 136-137, 139, 140-145, 146
- of ovary in, 138, 139, 140-145, 150-151
- of stomach in, 138, 140-145
- for carcinoma of breast in, 137-138, 140-145, 146-147, 148
- for other tumors in, 138, 140-145, 147-151
- patients with far advanced metastatic cancer studied in, 131-132
- grading of performance status in, 132
- socioeconomic status of, 131-132
- previous studies of, 131
- schedule of drug administration in, 132
- steric properties of, 131
- summary of cases of tumor regression in, 140-145
- types of previous therapy given in, 140-145
- Food, temperature of, as etiological factor in cancer of esophagus, study of, 389-413
- 5-FU. *See* 5-Fluorouracil
- Gallbladder, carcinoma of, in dogs, report of, 1127-1130
- method of inserting carcinogenic pellet in, 1127
- pathological findings in, 1127-1130
- appearance of mucosa in, 1127-1130
- chronic cholecystitis in, 1127-1130
- size of tumor in, 1127-1130
- use of methylcholanthrene as test carcinogen in, 1127-1130
- time of development of neoplasm in, 1127-1130
- Ganglioneuroma(s), of adrenal medulla, containing cells morphologically identical to hilus cells (extraparenchymal Leydig cells), report of, 421-425
- Gastrointestinal tract. *See also specific names of organs*
- involvement of, by giant follicular lymphoblastoma, report of, 759-760
- malignant lymphomas of, in African children, pathological entity of, 275
- metastases to, from carcinoma of breast, adrenal steroid-induced, report of, 426-431
- multiple lymphomatous polyposis of, 249-257
- absence of terminal leukemic state in, 254, 256
- age and sex distribution of patients
- Gastrointestinal tract, multiple lymphomatous polyposis of—(Contd.)
- with, 254, 255, 256
- case reports, 250-254
- classification of cases reported in literature and present cases in, 250
- deep X-ray therapy for, 256
- extent of alimentary tract involvement in, 254, 255, 256
- histological classification of, 255
- indications for noncurative surgical intervention in, 256
- mucosal and submucosal sites of, 254
- appearance of mucosal thickening in, 254, 256
- of raised nodular patches ("Peyer's patches"), 254
- of sessile and pedunculated polyps in, 254, 256
- pathological findings in cases of, 251, 252, 253, 254
- prognosis in, 256
- survival rates in, 255, 256
- symptoms in, 250, 251, 252, 253, 254, 256
- terminal generalized disease in, 254, 256
- Genetics, possible component in stomach cancer, report of, 199-200
- studies on inheritance of cancer of uterine cervix, 179-184
- Genital(s). *See also specific names of organs*
- female, measurement methods and results in study of quantitative characteristics of exfoliated cells from, 437-454
- cell characteristics selected for measurement in, 437
- 5 population distributions in, 444-453
- correction for variations in staining in, 439-440
- distribution of correction factor required in, 440
- problems with Panpanicolaou method in, 439
- significance of depth of stain in, 440
- definition of measurable cell in, 438
- determination of extinction coefficient of nucleus in, 438
- grouping factors in, 440-443
- age as, 443
- collection methods as, 441
- cytological classification as, 441-442
- menstrual cycle as, 443-444
- menopausal status as, 442
- normal and abnormal smears used in, 438
- peaks of aberrant cells in, 449-453
- previous methods of instrument screening in, 437
- random sample of cells in, 437
- relationship between cell diameter and nucleus diameter in, 444-453
- between extinction coefficient and diameter of nucleus in, 444-454
- results in, 444-454
- sample cell diameter histogram in, 441
- nuclear cell histogram in, 443
- tests of reliability of procedure used in, 438
- use of vertical stage projection microphotometer in, 438
- variations of extinction coefficient of nucleus in, 438, 439
- suitability of quantitative cytological measurements for automatic prescreening of cells exfoliated from, 455-468
- analytic methods used in, 455-459
- appearance of curves in, 457-459
- comparison of cumulative distribution curves for cervical smears from several different cytological categories in, 457-459
- instrument performance in, 467-468
- use of Cytoanalyzer in, 467
- results for cervical scrapings
- Genital(s), female—(Contd.)
- from premenopausal and postmenopausal women in, 464-466
- comparison with results for cervical smear in, 464, 465-466, 467
- using cell to nucleus (C/N) ratio aberrancy criteria in, 465
- nucleus aberrancy criteria in, 464-465
- for vaginal aspirations from premenopausal and postmenopausal women in, 466-467
- comparison of results for cervical smear and cervical scraping in, 466-467
- with postmenopausal cervical smears in, 462-466
- aberrancy curves for, 463
- atrophic and nonatrophic smears in, 462-463
- cell to nucleus (C/N) ratios in, 463-464
- contour plots for, 462-463
- with premenopausal cervical smears in, 460-462
- definition of smear classifications in, 460
- nucleus diameter and extinction coefficient in, 460
- role of cell to nucleus (C/N) diameter in, 461-462
- statistical variation in aberrant cell population in, 460-461
- total quantity of stained nucleus material in, 461
- use of more than one criterion in, 462
- various values of cell diameters in, 461
- screening performance in, 459-468
- classification of smears in, 459-460
- selection of value for screening level in, 458-459
- male, spermatocytic seminoma of, report of, 788, 794
- primary cancer of, frequency of occurrence of second primary malignant tumors in women with, 232
- Genital tract, female, mesonephric tumors of, 711-725
- age of patients in, 713-723
- case reports, 713-719
- description of mesonephric apparatus in, 711
- histological findings in, 712-723
- patients' history in, 713-723
- site of tumor in, 713-723
- size of tumor in, 713-719
- staining procedures used in, 713
- structure of tumor cells in, 711-725
- subgrouping of tumors in, 719-725
- tubular structure of tumors in, 711-725
- Genitourinary tract. *See also specific names of organs*
- involvement of, by giant follicular lymphoblastoma, report of, 763
- Geography, as etiological factor in cancer of esophagus, study of, 389-413
- GH. *See* Growth hormone
- GHADIALY, FERDZ N.: The role of the hair follicle in the origin and evolution of some cutaneous neoplasms of man and experimental animals, 801-816
- GILLIAM, ALEXANDER G.; MILLMORE, BENNO K., and LLOYD, J. WILLIAM: Trends of mortality attributed to carcinoma of the lung. The declining rate of increase, 622-628
- Gland(s). *See specific names*
- Glutathione, effect of, on radiation sensitivity of mouse tumors, 118, 119, 120
- Golter(s), solitary nodular and multinodular, removed at operation, differing frequencies of thyroid carcinoma in, explanation for, 67-71
- GOLDBERG, GIL M.; RUBENSTONE, ALBERT I., and SAPHIR, OTTO: A study of malignant lymphomas and leukemias. IV. Diagnosis of controversial cases of leukemia (with reference to histological criteria), 30-35
- RUBENSTONE, ALBERT I., and SAPHIR,

- OTTO: A study of malignant lymphomas and leukemias. III. Stem cell, blast cell, and monocytic leukemias (with reference to their lymphogenous or myelogenous origin), 21-29
- GOLDENBERG, IRA S., and HAYES, MARK A.: Hormonal therapy of metastatic female breast carcinoma. II. 2-Methyl dihydrotestosterone propionate, 705-706
- GONAD(S), malignant lymphomas of, in African children, pathological entity of, 275, 276
- GOODNER, JOHN T.; BERG, JOHN W., and WATSON, WILLIAM L.: The nonbenign nature of bronchial carcinoids and cylindromas, 539-546
- GORLIN, ROBERT J.; CHAUDHRY, ANAND P., and PRINDLES, JENS J.: Odontogenic tumors. Classification, histopathology, and clinical behavior in man and domesticated animals, 73-101
- GOSLING, J. R. G.; ABELL, M. R.; DROLETTE, BERNADETTE M., and LOUGHIN, T. D.: Infiltrative squamous cell (epidermoid) carcinoma of vulva, 330-343
- GT. *See* Glutamic oxalacetic transaminase
- GPT. *See* Glutamic pyruvic transaminase
- GR. *See* Glutathione reductase
- GRAY, EDWARD B., JR., *jt. au.*, 1122-1126
- GRECO, TOMMASO, *jt. au.*, [4]
- GRIFFITHS, JOHN D.; HOPPE, EVERETT, and COLE, WARREN H.: The influence of thermal stress and changes in body temperature on the development of carcinosarcoma 256 Walker in rats after inoculation of cells, 111-116
- GROSSI, C. E.; AGOSTINO, D.; MELAMED, M., and CLIFFORD, E. E.: The effect of human fibrinolytic on survival of carcinosarcoma 256 Walker cells in the blood, 957-962
- jt. au.*, 414-420
- GROVES, LAURENCE K., *jt. au.*, 1139-1143
- Growth, bone, effect of skeletal lesions of acute leukemia on, 608-621
- hormone (GH), effect of, on transplanted tumor in mice, 507-511
- in normal and malignant cells, quantitative assessment of alkaline phosphatase activity in, 502-506
- tumor, biomathematical approach to study of, 1272-1294
- H³. *See* Tritium
- HADDEN, D. H., *jt. au.*, 1057-1059
- Hair, follicles of, role of, in origin and evolution of benign epithelial tumors, report of, 801-816
- Hamartoma(s), bronchial, possible histogenetic relationship of bronchial carcinoids and salivary gland type tumors and, 169, 175
- Hamster(s), golden (*Mesocricetus auratus*), in study of growth curves of human cancer transplants during experimental chemotherapy with Methotrexate, report of, 1117-1121
- Hanks's balanced salt solution, in preparation of cultures of teratocarcinomas used in study of *in vitro* and *in vivo* differentiation, report of, 1017
- HARTMANN, WILLIAM H., and SHERLOCK, PAUL: Gastrointestinal metastases from carcinoma of the breast. An adrenal steroid-induced phenomenon, 426-431
- HAYASHI, Y.; COWDRY, E. V., and SUNTZEFF, V.: Microscopic properties of the basement membrane and elastic fibers of trachea and bronchus of smokers and nonsmokers, 1175-1182
- HAYES, DONALD M.: The seasonal incidence of acute leukemia. A contribution to the epidemiology of the disease, 1301-1305
- HAYES, MARK A., *jt. au.*, 705-706
- HAZARD, JOHN B., *jt. au.*, 1139-1143
- Heart, involvement of, by giant follicular lymphoblastoma, report of, 759
- Heat, as physical agent affecting radiation sensitivity of mouse tumors, report of, 121
- HEINRECKER, PETER, *jt. au.*, 862-874
- HELSPER, JAMES T., *jt. au.*, 512-516
- Hemangioblastoma(s), of cerebellum, report of, 784
- Hemangioendothelioma(s), malignant, in infants and children, report of, 1186-1196
- age in reported cases of, 1186-1188
- case reports, 1193-1195
- characteristic growth features in, 1189-1195
- clinical features of, 1188
- definition of tumor in, 1186
- differential diagnosis in, 1193
- hemangioendothelioma and, 1193
- hemangiopericytoma and, 1193
- so-called alveolar rhabdomyosarcoma and, 1193
- follow-up of patients in, 1193
- incidence of metastases in reported cases of, 1186-1188, 1195
- of recurrence of tumors in, 1188, 1192-1195
- location of reported tumors in, 1186-1188
- pathological features of tumors in, 1188-1189
- amount of collagen in, 1191
- anastomosing capillaries in, 1189
- appearance of endothelial cells in, 1189
- comparison of benign juvenile type and malignant type in, 1189
- illustration of malignant metastasizing type in, 1189-1190
- mitoses in, 1189-1190
- papillary proliferation in, 1190
- pattern of budding or sprouting of tumor cells in, 1190
- size of tumors in, 1189
- vascular clefts in, 1191
- reported incidence of different types of tumors in, 1186-1188
- sex of patients in, 1186-1188
- Hamangiopericytoma(s), relationship of, to glomus tumors in children, report of, 563-564
- Hepatotomy, middle lobe, for cancer of liver, report of, 1295-1300
- partial, for nutritional cirrhosis, autoradiographic and histological study of repair after, 963-967
- Hepatitis, activity of triphosphate isomerase in blood serum from patients with, 1166-1174
- Hepatoma(s), in mice, after exposure to cigarette tar and beta radiation, incidence of, 348, 353
- malignant, in infancy, complete right hepatic lobectomy for, 361-368
- Heredity, as factor in cancer of esophagus, study of, 392
- in multiple congenital tumors of soft tissues, possibility of, 913-920
- as possible factor in stomach cancer, analysis of 5 "stomach cancer families" in State of Utah to determine, 1005-1016
- of cancer of human uterine cervix, studies of, 179-184
- role of, in families of patients with multiple primary malignant tumors, report of, 234-235
- 5-HIAA. *See* 5-Hydroxyindoleacetic acid
- HILARIS, BASIL B., *jt. au.*, 577-581
- HINERMAN, DORIN L., *jt. au.*, 944-952
- Histiocyte(s), and histiocytic reaction, in cytological diagnosis of carcinoma of vagina, incidence of false positive risks in, 1223-1225
- Histiocytoma(s), in children, report of, 469-482
- case reports, 481
- clinical features of, 477-479, 480
- age at onset in, 477, 480
- anatomical location of tumor in, 478, 480
- average duration of tumor in, 477-478, 480
- confusion in nomenclature in, 469, 470
- differential diagnosis in, 479
- differentiation between fibrous xanthomas and, 470
- follow-up in, 479-480
- incidence of malignant tumors in, 479-480, 481
- of metastases in, 479-480, 481
- number of mitoses in tumors in, 479, 480
- pathological findings in, 479
- after gross examination in, 479
- after microscopic examination in, 478, 479
- Histogram(s), use of, in measurement study of quantitative characteristics of cells exfoliated from female genital tract, report of, 441, 444
- Histology, use of giant sections in study of spread of cancer of uterine cervix, report of, 42-54
- HITZELBERGER, WILLIAM E.; KERNOHAN, JAMES W., and UHLEIN, ALFRED: Giant cell fibrosarcoma of the brain, 841-852
- HN2. *See* Nitrogen mustard
- HOCHMAN, A., *jt. au.*, 300-307
- Hodgkin's disease, and lymphosarcoma, delayed skin response in, 597-607
- age of patients in, 599
- antigens used in, 599
- composition of study groups in, 598
- "definite" category in, 598
- "no or slight" category in, 598
- distribution of skin test responses to intermediate PPD and second strength PPD, *Trichophyton* extract, and *Candida* extract in, 600
- to mumps, tuberculin, *Trichophyton*, and *Candida* antigens among patients with Hodgkin's disease in, 603-604
- incidence of positive skin reactions among study groups, according to sex, in, 599
- of total energy to antigens used in, 600, 601
- possible causes of delayed skin response in cases of Hodgkin's disease in, 605-606
- race of patients in, 599
- responses to BCG vaccination among patients with lymphoma and healthy controls in, 602
- serial studies of tuberculin response in patients with lymphoma and leukemia in, 601
- sex of patients in, 599
- use of bacille Calmette Guérin (BCG) in, 598
- of rank t test in, 600, 601
- vaccination method used in, 599
- presenting as multiple polyposis of gastrointestinal tract, 249-257
- primary, of lung, report of, 1151-1165
- HOFFMANN, DIETRICH, *jt. au.*, 1306-1315
- HOLLAND, J. F., *jt. au.*, 205-209
- Homotransplantation, in addition to autotransplantation, in patients with advanced cancer, quantitative studies of, 971-978
- HOPPE, EVERETT, *jt. au.*, 111-116
- Hormone(s). *See also specific names of agents*
- and adrenalectomy, as therapy for advanced breast carcinoma, evaluation of, 1259-1265
- androsterone as, effect of, on clinical course and hormonal excretion in cancer of breast, 195-198
- effect of, on radiation sensitivity of mouse tumors, 120-121
- excretion of, effect of androstenediol dipropionate (5 α -androstan-3 α ,17 β -diol) in treatment of breast cancer on, 707-710
- role of, in luteinization of stroma in metastatic ovarian cancer, 827-840
- therapy with, and survival of patients with "recurrent" or inoperable breast carcinoma, 300-307
- use of 2 α -methyl dihydrotestosterone propionate for metastatic carcinoma of female breast, 705-706
- HOHN, ROBERT C., JR., *jt. au.*, 779-787
- HORVATH, W. J., *jt. au.*, 437-454, 455-468
- Host defense, in squamous cell carcinoma of esophagus, role of, 927-928
- HOYE, ROBERT C., and SMITH, ROBERT R.: The effectiveness of small amounts of preoperative irradiation in preventing the growth of tumor cells disseminated at surgery. An experimental study, 284-295
- HRESCHSHVYN, M. M.; SHERMAN, F. R., and HOLLAND, J. F.: Visualization of retroperitoneal lymph nodes. Lymphangiography as an aid in the measurement of tumor growth, 205-209
- 5-Ht. *See* 5-Hydroxytryptamine
- HUME, ROXANE, *jt. au.*, 883-888
- HURST, DORSEY W., and MEYER, OVID

- O.: Giant follicular lymphoblastoma, 753-778
- Hydrocarbon(s), polycyclic, as carcinogenic chemicals, in newborn mice, report of, 308-317
- 17-Hydroxycorticoid(s), plasma, total, response, in rats inoculated with carcinosarcoma 256 Walker cells, after thermal stress and body temperature changes, study of, 111-116
- 5-Hydroxyindoleacetic acid (5-HIAA), urinary, in adenomatoid bronchial tumors, levels of, 172-173
- 5-Hydroxytryptamine (5-HT), in adenomatoid bronchial tumors, role of, 171
- Hygiene, oral, as etiological factor in cancer of esophagus, study of, 389-413
- Hypercitricemia, in advanced cancer, and parathyroid function, relationship of, 934-943
- Hyperplasia(s), adrenocortical, in cases of bronchogenic carcinoma, previous studies of, 105, 107
- follicular, and follicular lymphoblastoma, comparison of, 770
- ovarian cortical stromal, from advanced cancer of breast in relation to therapeutic oophorectomy, significance of, 126-130
- Hypothermia, cancericidal effects of X irradiation and, 1061-1062
- Icn. See Isocitric dehydrogenase
- Ileum, as location of myxoma in children, report of, 634
- as site of carcinoid tumors, comparison of incidence of with that for small intestine, report of, 902-903
- Immunology. See Host defense
- Implantation; of methylcholanthrene as test carcinogen, in dogs, development of carcinoma of gallbladder after, 1127-1130
- Incidence, of leukemia, seasonal, epidemiological study of, 1301-1305
- INCE, JOSEPH S., *jt. aw.*, 105-110
- Infarct(s), bone, in acute leukemia, report of, 614
- Infarction, as possible etiological factor in peripheral lung cancer arising in scars, report of, 1213
- Infection(s). See also specific names
- and seasonal incidence of acute leukemia, possible relationship between, 1302-1305
- Infiltrate(s), stromal inflammatory cell, as prognostic factor in squamous cell carcinoma of esophagus, 921-933
- INGRAM, MARGARET, *jt. aw.*, 161-178
- Intestine(s), small, canine, response to irradiation, influence of succinylsulfathiazole (Sulfasuxidine), report of, 862-874
- comparison of Sulfasuxidine-treated and non-Sulfasuxidine-treated dogs in, 862-874
- findings by arterial angiography of mesentery in, 866-867
- disappearance of patent arteries and veins seen in, 866, 867
- gross appearance of ileum in, 863
- adhesions of omentum to ileum segments in, 863
- appearance of mesentery in, 863
- peristalsis in, 863
- lethality in, 863
- lymphangiographic findings in, 864-866
- mechanisms of obstruction of irradiated intestine in, 870-874
- microscopic appearance in, 866-874
- anastomosis of proximal and distal nonirradiated intestine in, 867-874
- development of granulation tissue in, 867-874
- disappearance of mucosa in, 866-874
- size of mucosally denuded area in, 867-874
- mongrel male dogs in, 862-874
- physical factors for radiation used in, 862
- size of ulcers in, 863-864
- method of plotting length of ulcers against time lapsing after irradiation in, report of, 863-864
- life history of carcinoid tumor of, 901-912
- Intestine(s), small—(Contd.)
- case reports, 907, 910
- clinical diagnosis in, 906, 910
- data for functioning carcinoid tumor in, 908-910
- average duration of symptoms in, 907, 909
- clinical manifestations of functioning carcinoid syndrome in, 908-909
- incidence of carcinoid syndrome in patients with carcinoid tumors of small intestine in, 910
- production and secretion of serotonin in, 908
- deep mucosal site of, 902
- duration of symptoms prior to diagnosis in, 907
- frequency of associated other primary malignant neoplasms and, 902-903, 905
- frequency of occurrence of, 901-902
- incidence by site of origin in, 902
- small intestine compared to ileum as, 902-903
- length of life prior to diagnosis in, 901
- method of spread of primary tumor in, 903
- natural history of carcinoid tumor in, 910-911
- slow rate of growth and metastasis in, 910-911
- pathological features of, 902-905
- roentgenological diagnosis in, 905, 909
- sex incidence in, 901
- signs and symptoms in, 905, 906
- abdominal pain as, 905, 906
- syndrome of intermittent obstruction of small intestine as, 906, 908
- size of primary tumor in, 902, 904
- surgery for, 906-907, 910
- 5-year survival rates, 906, 910
- tendency toward multicentricity of origin of, 902, 904
- IOB, VIVIAN; McMATH, MADELINE; COON, WILLIAM W., and BLOCK, GEORGE E.: Alterations in serum proteins of patients receiving palliative therapy for metastatic carcinoma of the breast, 1081-1084
- Irradiation, as therapy for giant follicular lymphoblastoma, report of, 770-771, 772-773
- preoperative, effectiveness in prevention of growth of tumor cells disseminated during surgery, experimental study of, 284-295
- IRVINE, E. S., *jt. aw.*, 636-638
- IRVINE, E. W., JR.; IRVINE, E. S., and MOFFITT, O. P., JR.: Treatment of laryngeal papillomas with bovine wart vaccine. A preliminary report of 4 cases, 636-638
- ISAACSON, C., *jt. aw.*, 1088-1089
- ISAACSON, E. ARNOLD, *jt. aw.*, 1005-1016
- Isotope(s), radioactive. See specific names
- Jaundice, obstructive, activity of triphosphate isomerase in blood serum from patients with, 1166-1174
- Jaw(s), as location of myxoma of children, report of, 630, 631-633
- malignant lymphomas of, in African children, clinical syndrome of, 262, 263-264
- pathological entity of, 270-272, 273
- upper, as location of congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, prognoma, and pigmented epulis of infancy, report of, 1090-1108
- JOHNSON, HORTON A., and BOND, VICTOR P.: A method of labeling tissues with tritiated thymidine in vitro and its use in comparing rates of cell proliferation in duct epithelium, fibroadenoma, and carcinoma of human breast, 639-643
- JONES, JOHN C., *jt. aw.*, 1151-1165
- KARBER, K., *jt. aw.*, 1197-1204
- KAUFFMAN, SHIRLEY L., and STOUT, ARTHUR PURDY: Histiocytic tumors (fibrous xanthoma and histiocytoma) in children, 469-482
- and STOUT, ARTHUR PURDY: Malignant hemangioendothelioma in infants and children, 1186-1196
- VON KAULLA, KURT N., *jt. aw.*, 889, 896
- KELLEY, RICHARD R., and SCULLY, ROBERT E.: Cancer developing in dermoid cysts of the ovary. A report of 8 cases, including a carcinoid and a leiomyosarcoma, 989-1000
- KELLY, J. E., *jt. aw.*, 131-152
- Keratocanthoma(s), berry-shaped, development of, after cutaneous carcinogenic exposure, role of hair follicle in, 801-816
- bud-shaped, development of, after cutaneous carcinogenic exposure, role of hair follicle in, 801-816
- dome-shaped, development of, after cutaneous carcinogenic exposure, role of hair follicle in, 801-816
- KERN, WILLIAM H.; CREPEAU, ANDRE G., and JOHN C.: Primary Hodgkin's disease of the lung. Report of 4 cases and review of the literature, 1151-1165
- KERNOHAN, JAMES W., *jt. aw.*, 841-852
- KETCHAM, ALFRED S.; KINSEY, DAVID L.; WEXLER, HILDA, and MARTEL, NATHAAN: The development of spontaneous metastases after the removal of a "primary" tumor. II. Standardization protocol of 5 animal tumors, 875-882
- jt. aw.*, 507-511
- 17-Ketosteroids (17-ks), urinary, after administration of androsterone in treatment of advanced cancer of breast, levels of, 196, 197, 198
- Kidney(s), carcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 244
- malignant lymphomas of, in African children, pathological entity of, 274
- mouse, infected with SE polyoma virus, difference in host cell-virus relationship between tubular epithelium and stroma in, 567-576
- cytochemical findings in, 570-571, 573
- characteristics of inclusions in, 570-571, 576
- deoxyribonucleic acid (DNA) behavior in cells in, 570, 573, 575
- cytopathological and histopathological findings in, 568-570
- "cellular activity" in, 568-570
- stromal hyperplasia in, 570, 572
- time sequence for appearance of changes in, 570
- "viral activity" in, 568-570
- degenerative and proliferative processes in, 571-575
- first cycle (intimate virus-host cell relationship) in, 574-575
- second cycle (absence of intimate relationship) in, 574-575
- histological and cytochemical methods used in, 568
- inoculation procedures in, 568
- origin of virus in, 567-568
- preparations of cell cultures in, 568
- use of NIH white mice in, 568
- KINSEY, DAVID L., *jt. aw.*, 875-882
- KRAUER, WARREN H., *jt. aw.*, 361-368
- KOPMAN, SYDNEY, *jt. aw.*, 644-648
- KOHOUT, ELYRIEDE, and STOUT, ARTHUR PURDY: The glomus tumor in children, 555-566
- KOSS, LEOPOLD G.; MELAMED, MYRON R., and DANIEL, WILLIAM W.: In situ epidermoid carcinoma of the cervix and vagina following radiotherapy for cervical cancer, 353-360
- KOTOB, NASSOUR, *jt. aw.*, 934-943
- KRAB. See Potassium β -aminobenzoate
- KRAMER, STANLEY P., and SELIGMAN, ARNOLD M.: The distribution of naphthylchloracetamidase with implications for the chemotherapy of cancer, 296-299
- LAKESSO, ANDREW O., *jt. aw.*, 913-920
- LAVIER, C. JAMES, and HINERMAN, DONALD L.: A morphologic study of pancreatic carcinoma with reference to multiple thrombi, 944-952

- LAGAK, F., *jt. sw.*, 1166-1174
 LANGLEY, F. A., *jt. sw.*, 711-725
 Large bowel, adenocarcinomas of, treated by 5-fluorouracil (5-FU), clinical evaluation of, 136-137, 139, 140-145, 146
 Laryngectomy, primary en bloc, and radical neck dissection for treatment of primary epidermoid carcinoma of larynx, incidence of cervical lymph nodes found in, 55-66
 subtotal, use of, in treatment of cancer of larynx, 1131-1138
 Larynx, cancer of, anatomical studies related to dissemination of, 1131-1138
 classification system used in, 1132
 clinical report in, 1135-1136
 need for numerous frozen section studies in, 1136
 survival rates in, 1135-1136
 technique of subtotal laryngectomy in, 1135
 demonstration of submucosal compartments in, 1132
 distribution of dye in, 1133
 "intrathyroid cartilage" in, 1132
 lack of communication between compartments in, 1132-1133
 factors affecting spread in, 1132
 findings for free margin of true vocal cord in, 1133-1134
 for lymphatics in, 1134-1135
 incidence of contralateral metastases in, 1134-1135
 prediction of route of spread in, 1131
 preliminary studies of, 1131
 types of cancer spread seen in, 1131
 types of metastatic involvement in, 1135
 carcinoma of, epidermoid, incidence of cervical lymph node metastases from and their relationship to certain characteristics of primary tumor, report of, 55-66
 age of patients and topographical distribution in, 56, 57
 bilateral and contralateral metastases in, 64
 cellular differentiation in, 57, 58
 cervical lymph node metastases in, 57, 58-59
 correlations with cervical metastases in, 59-62
 differentiation of tumor and, 58, 59
 distribution of tumors by site and type of peripheral tumor growth pattern in, 59-61
 fixation of vocal cord and intrinsic muscle invasion in, 59, 62
 incidence of nerve sheath invasion and, 58, 59
 invasion of small venous channels and, 59
 division of radical neck dissection specimens by region in, 57
 schematic drawing of, 57
 homogeneity of series in, 55, 56
 incidence of secondary radical neck dissection in, 56
 lymph node metastases and peripheral tumor growth pattern in, 58, 61-62
 metastatic rates in, 62-63
 comparison in patients with elective and therapeutic dissections in, 62, 63
 rate of clinically inapparent lymph node metastases in, 62, 63
 methods used in analysis of, 62-63
 accuracy in preoperative assessment of metastases in, 62-63
 size and site of tumor in, 56, 57
 subdivision of groups by anatomical sites of involvement in, 56-57
 use of patients treated by primary en bloc laryngectomy and radical neck dissection in, 55
 epitheliomas of, as multiple primary malignant tumors of multicentric origin, report of, 239
 papillomas of, treatment of, with bovine wart vaccine, preliminary report of, 636-638
 case reports, 636-638
 Larynx, papillomas of—(Contd.)
 clinical trial in, 636
 doses of bovine wart vaccine used in, 636-638
 lack of recurrence after treatment with, 636-638
 possible viral agent involved in, 638
 previous operative procedures in patients with, 636-638
 symptoms in patients with, 636-638
 therapy with drugs in patients with, 636-638
 LD₅₀. See Median lethal dose
 LEFFALL, LA SALLE D., *jt. sw.*, 1127-1130
 Leiomyoma(s), uterine, leiomyosarcoma in situ and, 483-492
 Leiomyosarcoma(s), in dermoid cysts of ovary, report of, 989-999
 LEMON, HENRY M., and KOTOK, NASSOUR: Hypertrichemia and parathyroid function in advanced cancer, 934-943
 Lesion(s), bronchial, dysplastic, histological study of, 1226-1234
 osteolytic, in acute leukemia, 614
 renal pelvic, exfoliative cytology of, 1085-1087
 satellite, differentiation from primary tumor growth, biomathematical approach to, 1289-1293
 Leucovorin. See Citrovorum factor
 LEUCHTENBERGER, CECILIE, *jt. sw.*, 567-576
 LEUCHTENBERGER, RUDOLF; LEUCHTENBERGER, CECILIE; STEWART, SARAH E., and EDDY, BERNICE E.: Difference in host cell-virus relationship between tubular epithelium and stroma in kidneys of mice infected with SE polyoma virus. A correlated cytological, histological, and cytochemical study, 567-576
 Leukemia(s), absence of, in cases of malignant lymphoma in African children, report of, 282
 activity of triphosphate isomerase in blood serum from patients with, 1066-1074
 acute, seasonal incidence of, epidemiological study of, 1301-1305
 month of onset of symptoms and month of histological diagnosis in, 1301, 1302, 1303
 possible association of endocrine organ weight variation in laboratory animals and leukemia incidence curve in, 1303, 1305
 relationship between infectious diseases and, 1302-1305
 statistical comparison with hypothetical random distribution in, 1303-1305
 skeletal lesions of, 608-621
 age and sex of patients in, 608
 alteration in endochondral bone formation in, 618-619
 association of bone pain with roentgenological evidence of bone lesions in, 612-613
 evolution of bone lesions in, 613
 classification of patients by type of leukemia in, 608
 clinical findings in, 608-610
 fever associated with, 609
 joint involvement in, 609
 cortical and periosteal lesions in, 612
 histological sections used in study of, 608
 pathological findings in, 613-616
 bone infarcts in, 614
 "growth arrest" lines in, 615
 knee and hip joint involvement in, 616
 leukemic infiltration and masses in periosteum in, 614-615
 osteitis fibrosa in, 614
 osteolytic lesions in, 614
 osteoporosis in, 613-614
 radiolucent bands in, 615-616
 suppurative arthritis due to *Pseudomonas aeruginosa* in, 616
 patient growth rate in, 609, 610
 duration of disease in relation to, 609, 610
 Leukemia, acute, seasonal incidence of—(Contd.)
 roentgenographic examinations on admission in, 608, 610-613
 osteolytic lesions in, 611-612
 radiolucent bands seen in, 610, 611
 relative incidence, type, and distribution of bone lesions in, 610, 611
 variety of lesions found in, 610
 treatment with colchicine in, 609-610
 blast cell, in relation to lymphogenous or myelogenous origin, study of, 21-29
 clinical data for patients with, 22-23
 diagnosis of, according to conventional criteria, in, 23
 according to lymphatic involvement, in, 23
 histological appearance of, 21-24, 26-29
 diagnosis of controversial cases of, 30-35
 changes in pattern of tissue involvement in, 34
 data for cases in, 30-31
 diagnoses by conventional criteria in, 30-31
 final diagnoses in, 31, 34
 role of involvement of lymphatics in, 31-35
 staining methods used in, 31
 in mice, after exposure to cigarette tar and beta radiation, incidence of, 346, 347-348
 failure to demonstrate acceleration of, by cell-free brain filtrates from leukemic animals of same strain, report of, 496-501
 age at death for first mouse with leukemia in each group in, 499
 of animals at time of injection in, 497
 causes of deaths of animals in, 499, 500
 incidence of polyoma virus infection in, 499-501
 injection volumes in, 497
 leukemia incidence in experimental and control animals in, 498-501
 preparation of mouse brain filtrates in, 497
 serological studies carried out in, 498
 sources of animal loss from the experiments in, 497-498
 of mice in, 496
 "inbred" mice in, 496
 "penbred" mice in, 497
 survival times for experimental and control animals in, 498-499
 use of *Escherichia coli* organisms in, 497
 incidence of, in Singapore children, report of, 535, 536-537
 monocytic, in relation to lymphogenous or myelogenous origin, study of, 21-29
 clinical data for patients with, 22-23
 mononuclear, diagnosis of, by preponderant cell type in, 21-27
 data for cases studied in, 22-23
 stem cell, in relation to lymphogenous or myelogenous origin, study of, 21-29
 clinical data for patients with, 22-23
 Leukeran (chlorambucil), in treatment of childhood cancer, use of, 36-41
 Leukoplakia, presence and incidence of, in multiple oral carcinomas, report of, 513-514, 515
 LEWIS, F. JOHN, *jt. sw.*, 1060-1062
 Lip(s), epitheliomas of, as multiple primary malignant tumors of multicentric origin, report of, 239
 Liposarcoma(s), in children, as part of malignant compound tumor, report of, 524-533
 Liver. See also Cirrhosis
 cancer of, middle hepatic lobec-

Liver, cancer of—(Contd.)

- tomy for, 1295-1300
- case reports, 1297-1299
- indications for middle lobe hepatectomy in, 1295-1297
- patterns of anatomical arrangements of middle hepatic vein in relation to, 1296
- operative procedures in, 1297-1299
- present concept of classification of hepatic lobes in relation to, 1295-1296
- nomenclature of hepatic lobes and segments in relation to, 1295, 1296
- technique of middle hepatic lobectomy (median left segmentectomy) in, 1299-1300
- teratoid, in infancy, complete right hepatic lobectomy for, 361-368
- case report, 361-363
- etiology of, 363-364
- incidence rates for, 365
- normal fasting sugar level in, 366
- blood volume studies in, 367
- operative procedure in, 362
- pathological findings in, 362-363
- gross examination of specimen in, 362-363, 364
- microscopic examination of specimen in, 363
- postoperative care in, 366
- presence of anemia in, 366
- relationship of liver anatomy to, 365-366
- control of hemorrhage in, 366
- of hemostasis in, 366
- ligation of right portal vein in, 366
- role of falciform ligament in, 365
- spasm of vena cava in, 366
- step-by-step procedure suggested for dissection in, 365-366
- variations in hepatic pedicle in, 365
- of hemogram in, 366
- examination of, in differential diagnosis of leukemia, use of, 26-27
- hepatomas of, as multiple primary malignant tumors of multicentric origin, report of, 244
- involvement of, by giant follicular lymphoblastoma, report of, 761
- malignant lymphomas of, in African children, pathological entity of, 273-274
- metastases, experimental studies of effect of nutrition on, 547-554
- alteration of "soil" of liver in, 549-553
- composition of synthetic diets used in, 548
- effect of altering content of fat in diet in, 550-553
- of altering diet caloric intake in, 549-553
- of altering protein content of diet in, 549-550, 551-553
- of choline levels in, 550-553
- on tumor "take" in, 549-553
- relationship to evaluation of chemotherapeutic agents, hormones, irradiation, and other forms of therapy in, 553
- use of adult female Sprague-Dawley rats in, 547
- of carcinosarcoma 256 Walker in, 548
- metastatic involvement of, in patients with generalized cancer, activity of triphosphate isomerase in blood serum from, 1166-1174
- nutritional cirrhosis of, autoradiographic and histological study of repair after partial hepatectomy in, 963-970
- LLOYD, J. WILLIAM, Jr. *et al.*, 622-628
- Lobectomy, for primary Hodgkin's disease of lung, results of, 1151-1165
- hepatic, right, complete, for teratoid cancer of liver in infancy, report of, 361-368
- middle hepatic, for liver cancer, report of, 1295-1300
- LOCK, D. R. *et al.*, 1226-1234
- LOUGHRAN, T. D., *et al.*, 330-343
- Lower respiratory tract. *See also specific names of organs*
- involvement of, by giant follicular

Lower respiratory tract—(Contd.)

- lymphoblastoma, report of, 759
- LS402A VI tumor, in vitro and in vivo study of differentiation of teratocarcinomas using, 1017-1029
- Lung(s), cancer of, peripheral, arising in scars, report of, 1205-1215
- age of patients in, 1211, 1213
- autopsy findings for patients included in, 1205-1215
- case reports, 1205-1211
- clinical symptoms in, 1205-1215
- diagnosis in, 1205, 1206, 1209, 1210, 1211-1212
- duration of tumors in, 1213
- etiology of scar in relation to development of tumors in, 1213-1214
- gross appearance of tumors in, 1205-1215
- histological characteristics of scar cancer in, 1214
- types of cancers in, 1214
- incidence of metastases in, 1205-1215
- location of tumors in, 1205-1215
- microscopic appearance of tumors in, 1205-1215
- morphological variations in, 1215
- postoperative follow-up in, 1205-1215
- sex of patients in, 1211, 1213
- size of tumors in, 1205-1215
- symptoms in, 1205-1211, 1212
- types of metastases in, 1205-1215
- relationship of certain factors to survival of patients treated by pulmonary resection for, 1251-1258
- blood vessel invasion by cancer and survival rates in, 1253, 1255
- bronchial epithelial alterations and survival rates in, 1256-1257
- criteria for diagnosis of carcinoma *in situ* and, 1256
- cellular types of tumors in, 1252-1253
- adenocarcinoma as, 1252
- epidermoid carcinoma as, 1252
- undifferentiated carcinoma as, 1252
- clinical data in, 1251-1252
- types of resections used in patients in, 1251, 1252
- of symptoms in, 1251
- lymph node metastases and survival rates in, 1255-1256
- method of obtaining diagnosis in, 1251-1252
- microscopic findings associated with poor prognosis in, 1257-1258
- nerve sheath invasion and survival rates in, 1256
- pulmonary site of cancer and survival rates in, 1252-1253
- site and cell type of cancer and survival rates in, 1252-1253, 1257
- survival rates by type of operation in, 1252
- validity and value of histology and cytology classifications of, 673-698
- accuracy of cytological classification in, 687-688
- comparison of exfoliated cell types and histological types of tumor as alternatively classified in, 689
- correlation of cytological and histological appearances in, 688
- discrepancies between cytological and histological classification in, 688-689, 691
- average ages of patients in, 695-696
- and 5-year survival rates in, 673
- correlation of present findings with previous findings in, 691-695
- evaluation of alternative histological classifications in, 685-687
- histological diagnoses based on biopsy and corresponding operative specimens in, 685

Lung(s), cancer of—(Contd.)

- homogeneity of tumors and validity of morphological subdivisions in, 682-683
- pattern of tumor differentiation in, 683
- types of tissues seen singly or in combination in, 682, 683
- lung cancer categories in, 673-674
- original diagnoses in, 675
- relative frequencies of lung cancer in, 674
- second classification by most malignant component of tumor in, 686
- statistical treatment of data in, 679, 681
- equation for casualty rate in, 675, 679
- standard deviation estimates by tumor classifications in, 681
- use of life tables or statistical graphs in, 679, 681
- of modified median for calculation of average logarithmic duration in, 681
- survival times in, 695
- terminology in, 676-679
- validity of biopsy classification in, 683, 685
- value of original classification in indicating differences in survival times, sex incidence, and ages in, 681-682
- carcinoma of, adrenocortical width in, 105-110
- comparison of cell types in bronchogenic carcinoma group in, 108, 109
- of groups after exclusion of cortical hyperplasia in, 107
- comparison of groups in, 106-107
- groups of adrenal sections used in, 105
- diagnostic classification of, 106
- influence of microscopic adrenocortical metastases on thickness of adrenal cortex in bronchogenic carcinoma group in, 107-108
- technique of measuring cortical thickness in, 105-106
- total and zonal adrenal cortical thickness in bronchogenic carcinoma group by cell type in, 109
- cytological use of "wet films" in detection of, 1139-1143
- bronchoscopic specimens used in study of, 1139-1140
- cytological features found in, 1140
- difference between "wet film" method and "fixed film" method in, 1139
- false positive reports in, 1142
- "indeterminate" reports in, 1142
- results for adenocarcinoma in, 1140-1141
- for clinical carcinoma in, 1142
- for large cell type of undifferentiated carcinoma in, 1141
- for other pulmonary neoplasms in, 1142
- bronchial adenoma as, 1142
- malignant lymphoma as, 1142
- for secondary carcinoma in, 1142
- for small cell carcinoma in, 1141-1142
- for squamous cell carcinoma in, 1140
- staining solution used in, 1139
- time period covered in study of, 1139
- types of stains used in, 1139-1140
- giant cell, epithelial origin of confirmed by tissue culture, report of, 1052-1056
- case report, 1052-1054
- histological characteristics of, 1052, 1053, 1054-1055
- bizarre large cells in, 1052, 1053

- Lung(s) carcinoma, giant cell—(Contd.)**
evidence of phagocytosis in, 1052
stains used in, 1052
microscopic examination of tumor specimen in, 1053, 1054
pathological findings in, 1054
site of origin of tumors in, 1052
tissue culture procedure in, 1055
levels of amylase activity in, 1109-1114
trends of mortality attributed to, 622-628
age variations and changes in rates in, 624, 626, 627
unknown factors involved in predictions of, 627-628
annual percentage changes in rates in, 624, 625, 626-628
declining rate of increase found in, 625-628
definition of "cancer of lung" in, 622
mortality trends in, 622-625
age adjustment in, 623-625
problems in cause of death allocation in, 622
use of pancreatic deoxyribonuclease (dornase) in cytological diagnosis of, 419, 420
- Hodgkin's disease of, primary, report of, 1151-1165**
age of patients in, 1151-1161
case reports, 1151-1158
chemotherapeutic agents used in treatment of, 1152-1162
duration of symptoms in, 1151-1162
follow-up period in, 1152-1162
mediastinal lymph node involvement in, 1152-1162
methods of treatment used in, 1151-1165
microscopy of tumors, 1152-1164
race of patients in, 1151-1161
results of bronchoscopic examinations in, 1152-1162
of routine roentgenography in, 1151-1162
site of tumor in, 1152-1162
size of tumor in, 1152-1162
survival period after treatment in, 1152-1162
types of symptoms in, 1151-1162
metastases to, in mice, after removal of "primary" tumor, experimental study of, standardization protocol of 5 animal tumors in, 875-882
- Luteinization, of stroma, in metastatic ovarian cancer, endocrine significance of, 827-840**
- LURIE, H. I.: Congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, progonoma, and pigmented epulis of infancy. Summary and review of the literature and report of the first case in an adult, 1090-1108**
- and ISAACSON, C.: A melanotic progonoma in the scapula region, 1088-1089**
- Lymph node(s), bilateral dissection of, and complete vulvectomy, as treatment for Bowen's type of intraepithelial and infiltrative carcinoma of vulva, report of, 323-324, 328-329**
cervical, metastases to, from epidermoid carcinoma of larynx, incidence of and relationship to certain characteristics of primary tumor, report of, 55-66
from multiple oral carcinomas, report of, 514-515
dissection of, bilateral, and radical vulvectomy, as treatment for infiltrative squamous cell (epidermoid) vulvar carcinoma, results of, 333, 341-343
involvement of, in giant follicular lymphoblastoma, 755-757
in malignant lymphomas, in African children, 272-273, 280-281
mediastinal, metastases to, from primary Hodgkin's disease of lung, prognosis and, 1151-1165
metastases to, and wound-washing results, after operation for cancer, lack of correlation between, 1042, 1043
from cancer of vulva, 14-20
- Lymph node(s), bilateral dissection of—(Contd.)**
of cancer patients, excretion and distribution studies of radioactive labeled alkylating drugs in, 955, 956
retroperitoneal, lymphangiography as aid in measurement of tumor growth through visualization of, 205-209
case reports, 206-207
incision on dorsum of foot in, 205
injection of radio-opaque dye in, 205-206
radio-opacification of pelvic and para-aortic nodes after, 206
lymphangiograms taken after, 206-208
- Lymphangiosarcoma(s), developing at site of lymphedema after radical excision of breast cancer, incidence of, 233-234**
possible systemic carcinogenic influence in, 233-234
postmastectomy, in lymphedematous arm, treatment of, 210-212
case report, 210-211
early recognition and biopsy in, 211
indications for X-ray and/or surgical treatment in, 211-212
interscapulothoracic amputation and excision of local recurrence in, 210-211
survival after, 210-211
- Lymphatic system, involvement of, in blast cell leukemia, in relation to lymphogenous or myelogenous origin, study of, 21-29**
in stem cell leukemia, in relation to lymphogenous or myelogenous origin, study of, 21-29
diagnosis of, according to conventional criteria, in, 23
according to lymphatic involvement, in, 23, 26-28
histological appearance of, 22-23, 25, 26
- Lymphedema, of arm, postmastectomy, development of lymphangiosarcoma and, 210-212**
- Lymphoblastoma(s), giant follicular, report of, 753-778**
bone involvement in, 762-763
case reports, 755, 756, 757, 760, 761, 762-763, 764-765, 767, 771
chemotherapy with chlorambucil in, 771, 772-773
with nitrogen mustard (HN2) in, 771, 772-773
with triethylene melamine (TEM) in, 771, 772-773
classification in, 769
clinical features in, 754-763
course and prognosis in, 766-768
duration of disease in, 767-768
relation of symptoms to survival time and, 767-768
transitions to lymphosarcoma in, 768
differential diagnosis in, 769-770
comparison of follicular lymphoblastoma and follicular hyperplasia in, 770
eye involvement in, 757-758
fever in, 763
gastrointestinal tract involvement in, 759-760
genitourinary tract involvement in, 763
gross and microscopic findings in, 768-769
incidence of, 754
in relation to other lymphomas, 754
involvement of heart in, 759
of pericardium in, 759
irradiation for, 770-771, 772-773
laboratory findings in, 763-766
basal metabolic rate in, 766
for bone marrow, 765
for erythrocytes and hemoglobin, 763
for leukocytes, 763-765
for platelets, 765-766
for sedimentation rate, 766
for serum proteins, 766
for urine, 766
serological findings in, 766
liver involvement in, 761
lower respiratory tract tumors in, 759
- Lymphoblastoma(s), giant follicular—(Contd.)**
nervous system involvement in, 762
nonspecific lesions in, 757
organ or site involved in, 755
pancreatic involvement in, 762
parotid gland tumors in, 758
pathological findings in, 768-769
skin lesions in, 756-757
spleen involvement in, 761-762
surgical treatment in, 772-773, 774
thyroid tumors in, 758-759
treatment with cortisone and adrenocorticotrophic hormones in, 771, 772-773
types of transitions to other forms of lymphoma in, 769
presence of more than one type of lymphoma in, 769
upper respiratory tract tumors in, 758
weight loss in, 763
- Lymphoma(s), CD5, in experimental study of effectiveness of small amounts of preoperative irradiation in preventing growth of tumor cells disseminated at surgery, 284-295**
giant follicular, lymphoblastic, report of, 753-778
present as multiple polyposis of gastrointestinal tract, 249-257
Lewis, in Lewis strain adult rats, cancericidal effects of X irradiation at body temperature below 10° C. on, 1060-1062
malignant, and leukemias, study of, 21-29
in African children, clinical syndrome of, 258-269
abdominal tumors as, 264
differential diagnosis in, 264
histological confirmation in, 264
multiplicity or massiveness of lesions in, 264
classification of, 258-259
clinical presentation in, 262-266
course and therapy in, 266-267
4-amino-N¹⁰-methyl pteroylglutamic acid (Methotrexate) in, 266
radiosensitivity of tumor as factor in, 266
rapid growth rate of tumor in, 266
exclusion of Hodgkin's disease in study of, 258, 259
geographical distribution of, 260-261
incidence by presenting features in, 262
jaw tumors as, 262, 263-264
clinical involvement of mandible or maxilla in, 263
displacement of teeth in, 263
edema of eyelids in, 263
lack of skin surface ulceration in, 263
possibility of multicentric development in, 264
roentgenographic appearance in, 263, 264
site of origin of tumors in, 263-264
multiplicity of lesions in, 265-266
paraplegia as, 264-265
involvement of vertebral canal as cause of, 264-265
lack of radiological changes in bone in, 265
patient age and sex in, 261
possible environmental basis for high incidence and geographical distribution of, 267
manifestations of leukemia in, 268
presence of anemia in, 268
racial and tribal incidence in, 261-262
salivary gland tumors as, 265
skeletal (nonjaw) tumors as, 265

- Lymphoma(s), malignant—(Contd.)**
 pathological entity of, 270-283
 absence of leukemia occurring with, 282
 classification of tumors in, 277
 distribution by histological type of tumor in, 277
 histiocytic type in, 279, 281
 microscopic pathology in, 277-282
 mixed cell type in, 279
 organ distribution and gross pathology in, 270-277
 poorly differentiated lymphocytic type in, 277-278
 possible "priming" action of some parasite in, 282
 relationship of virus infection as "priming" factor and, 282
 principle anatomical sites of tumors in, 282
 abdominal viscera as, 273-275
 adrenal involvement in, 274-275
 gastrointestinal tract involvement in, 275
 kidney involvement in, 274
 liver involvement in, 273-274
 pancreatic involvement in, 275
 spleen involvement in, 275
 central nervous system as, 275-276
 paraplegia resulting from, 276
 gonads as, 275, 276
 involvement of ovaries in, 275, 276
 of testes in, 275
 jaw bones as, 270-272, 273
 lymph nodes as, 272-273, 280-281
 external lymphadenopathy as presenting symptom and, 273
 miscellaneous sites as, 276
 skeleton as, 272
 infiltrates in vertebrae in, 272
 use of bone marrow studies, 272
 of roentgenographs, 272
 thyroid and salivary glands as, 275
 role of possible primitive mesenchymal cell in, 281-282
 diagrammatic representation of malignant lymphoma histogenesis from, 281
 stem cell type in, 279, 280
 in mice after exposure to carcinogenic chemicals, 308-317
 lung, cytological use of "wet films" in detection of, 1142
 presenting as multiple polyposis of gastrointestinal tract, 249-257
- Lymphosarcoma(s), and Hodgkin's disease, delayed skin response in, 597-607**
 presenting as multiple polyposis of gastrointestinal tract, 249-257
 transitions from giant follicular lymphoblastoma to, 768
- MacDONALD, RICHARD A., *jt. au.*, 963-970**
 Macrophage(s), stromal, cytochemical study of esterase activity of, 369-377
- MAGNER, D., *jt. au.*, 817-826**
 Mammary duct system, involvement of, by Paget's disease, 653-672
- Mammary gland(s). *See* Breast(s)**
- Mandible(s), as location of congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, progonoma, and pigmented epulis of infancy, report of, 1090-1108**
 involvement of, in malignant lymphomas, in African children, incidence of, 262, 263-264
 malignant lymphomas of, in African children, pathological entity of, 270-272, 273
- MANTREL, NATHAN, *jt. au.*, 507-511, 875-882**
- MARSHALL, RICHARD B., and HORN, ROBERT C., JR.: Nonchromaffin paraganglioma. A comparative study, 779-787**
- Mastectomy, treatise on, by Angelo Nannoni, 14**
- Maxilla(e), as location of congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, progonoma, and pigmented epulis of infancy, report of, 1090-1108**
 involvement of, in malignant lymphomas, in African children, incidence of, 262, 263-264
 malignant lymphomas of, in African children, pathological entity of, 270-272, 273
- Mc (3-methylcholanthrene), as carcinogenic chemical, in newborn mice, report of, 308-317**
- MCCORMACK, LAWRENCE J.; HAZARD, JOHN B.; BELOVICH, DORIS; EPLYER, DONALD B., and GROVES, LAURENCE K.: Cytological use of "wet films" in the detection of carcinoma of the lung, 1139-1143**
- MCDONALD, G. O., *jt. au.*, 1057-1059**
- McGAVRAN, MALCOLM H.; BAUER, WALTER C., and OGURA, JOSEPH H.: The incidence of cervical lymph node metastases from epidermoid carcinoma of the larynx and their relationship to certain characteristics of the primary tumor. A study based on the clinical and pathological findings for 96 patients treated by primary en bloc aryngectomy and radical neck dissection, 55-66**
- McMATH, MADELINE, *jt. au.*, 1081-1084**
 Md. *See* Malic dehydrogenase
- Measurement, cytological, quantitative, suitability of, for automatic prescreening of exfoliated cells from female genital tract, report of, 455-468**
 interferometric studies of anhydrous mass of exfoliated cells from experimental cancer of cervix by, 378-383
 methods of, in study of quantitative characteristics of exfoliated cells from female genital tract, report of, 437-453
 of tumor growth, by use of lymphangiography of retroperitoneal lymph nodes, report of, 205-209
- Median lethal dose (LD₅₀), effect of hypothermia in rats on, 1061**
- MELAMED, MYRON R.; ROBBINS, GUY F., and FOOTE, FRANK W., JR.: Prognostic significance of gelatinous mammary carcinoma, 699-704**
jt. au., 353-360, 957-962
- Melanocarcinoma(s), congenital, melanotic adamantinoma, retinal anlage tumor, progonoma, and pigmented epulis of infancy, 1090-1108**
 case reports, 1094, 1097-1099
 histology of tumors in, 1090-1108
 mandible as location of, 1090-1108
 maxilla as location of, 1090-1108
 nose and oral cavity as locations of, 1103, 1106
 occurrence in adult of, 1097-1099
 opinions on histogenesis of, 1099-1107
 association of abnormal dental elements and, 1099-1103
 presence of melanin in, 1090-1108
 previous reports of, 1090-1097, 1099-1108
 skull as location of, 1090-1108
 upper jaw as location of, 1090-1108
- Melanoma(s), amelanotic, S91A, in study of spontaneous development of metastases after removal of "primary" tumor, use of, 875-882**
 juvenile, histochemical study of, 1001-1004
 case reports, 1001
 histopathological diagnosis in, 1001-1003
 alkaline phosphatase reactions in, 1003
 cholinesterase reactivity in, 1002, 1003
 dopa and tryrosinase reactions in, 1003
 epithelioid appearance in, 1002, 1003
 staining for polysaccharides and lipids in, 1003
- Melanoma(s), amelanotic—(Contd.)**
 use of hematoxylin and eosin-stained sections for 1001, 1002, 1003
 lack of pigmentation in, 1003
 vascularity of tumor in, 1003
- S91 Cloudman, in experimental study of spontaneous development of metastases after removal of "primary" tumor, use of, 875-882**
 in study of effectiveness of small amounts of preoperative irradiation in preventing growth of tumor cells disseminated at surgery, 284-295
- Membrane(s), basement, and elastic fibers of trachea and bronchus, of smokers and nonsmokers, microscopic properties of, 1175-1182**
- Menopause, as grouping factor in measurement study of cells exfoliated from female genital tract, 442**
 importance of, in study of suitability of quantitative cytological measurements for automatic prescreening of exfoliated cells from female genital tract, report on, 455-468
- Menstrual cycle, as grouping factor in measurement study of cells exfoliated from female genital tract, 442-443**
- MERANZE, DAVID R., *jt. au.*, 67-72**
- MERRILL, JAMES A., and ROSS, NEAL L.: Cancer of the vulva, 13-20**
- Mesenchymoma(s), malignant, in children, report of, 524-533**
 age of patient in, 525, 526, 527, 528
 at diagnosis, 525, 526, 527, 528
 at onset, 525, 526, 527, 528
 anatomical distribution of tumors in, 525, 526, 527, 528
 case reports, 529-532
 clinical features in, 528-533
 correlation of size of tumor and metastases in, 529
 and rate of recurrence in, 529
 and survival in, 529
 differentiated, and undifferentiated elements in, 527-533
 gross characteristics of tumors in, 525-527
 histological features of tumors in, 526-533
 incidence of metastases in, 526-533
 microscopic characteristics of tumors in, 526-533
 recurrence in, 526-533
 sex of patients in, 525, 526, 527, 528
 size of tumors in, 526-533
 survival in, 526-533
 treatment in, 526-533
- Metabolite(s)-antimetabolite cancer chemotherapy, using continuous intra-arterial 4-amino-N¹⁰-methyl pteroylglutamic acid (Methotrexate) with intermittent intramuscular citrovorum factor (CF), report of, 744-752**
- Metaplasia(s). *See* also Cell(s), Epithelium(s), and Nucleus(s)**
 of uterine cervix, relative maturation of surface epithelium in, 1045-1051
- Metastasis(es). *See* also specific sites**
 adrenocortical, microscopic, in bronchogenic carcinoma, influence of, on thickness of adrenal cortex, 107-108
 and multiple primary malignant tumors of multicentric origin, differential diagnosis of, 238-248
 axillary, from gelatinous breast carcinoma, significance of, 699-704
 cervical lymph node, from multiple oral carcinomas, involvement by, 514-515
 differentiation from primary tumor growth, biomathematical approach to, 1289-1293
 from adenoid cystic carcinoma, clinicopathological study of, 1235-1250
 from basal cell cancer, 1036-1040
 from Bowen's type of intraepithelial and infiltrative carcinoma of vulva, prognostic significance of, 318-329
 from breast cancer, effect of androsterone on clinical course and hormonal excretion on, 195-198
 effect of on survival rates, re-evaluation of, 1122-1126
 from breast carcinoma, alterations in serum protein levels in patients receiving palliative therapy for, 1081-1084

- Metastasis(es), from breast carcinoma—(Contd.)
 comparative evaluation of adrenalectomy and cortisone therapy on, 1259-1265
 from bronchial carcinoids and cylindromas, incidence of, 542, 543, 545, 546
 from cancer of larynx, anatomical studies in relation to dissemination in, 1131-1138
 of ovary, luteinization of stroma in, endocrine significance of, 827-840
 from carcinoid tumors of small intestine, 902, 904, 910
 from carcinoma of breast, levels of enzymes in serum of patients with, 726-733
 from carcinoma of pancreas, and multiple thrombi, relationship of, 948-949
 from female breast cancer, effect of androstenediol dipropionate (5 α -androstane-3 α ,17 β -diol) therapy on clinical course and hormonal excretion in, 707-710
 hormonal therapy with 2 α -methyl dihydrotestosterone propionate in, 705-706
 from lung cancer, relationship of, to survival of patients with pulmonary resections, 1251-1258
 from malignant hemangioendothelioma in children, incidence of, 1186-1196
 from mesenchymomas, in children, incidence of, 526-533
 from peripheral lung cancer arising in scars, report of, 1205-1215
 from primary Hodgkin's disease of lung, prognosis and, 1151-1165
 from "recurrent" or inoperable breast carcinoma, survival rates after hormonal treatment for, 300-307
 from thyroid carcinoma, in children, report of, 734-743
 from tumors developing in mice after exposure to cigarette tar and beta radiation, incidence of, 346, 347, 348, 350
 gastrointestinal, from carcinoma of breast, adrenal steroid-induced, report of, 426-431
 hepatic, experimental studies of effect of nutrition on, 547-554
 in adenomatoid bronchial tumors, role of, 170
 intracranial, value of radiation therapy in management of, 577-581
 liver, in patients with generalized cancer, activity of triphosphate isomerase in blood serum from, 1166-1174
 lung, in mice, after removal of "primary" tumor, experimental study of, standardization protocol of 5 animal tumors used in, 875-882
 lymph node, axillary, from breast cancer, significance of, 1266-1271
 to cervix, from epidermoid carcinoma of larynx, incidence of and relationship to certain characteristics of primary tumor, report of, 55-66
 occult, from breast cancer, significance of, 1266-1271
 pulmonary, time of development of, and release of circulating tumor cells, as determined in experimental study, report of, 883-888
 spontaneous, development of, after removal of "primary" tumor, standardization protocol of 5 animal tumors in, 875-882
 amelanotic melanoma S91A in DBA/p mice in, 875
 melanoma S91 Cloudman in thigh muscle of CDBA/2F₁ mice in, 875
 osteogenic sarcoma 112 in C3H/HeN mice in, 875
 sarcoma dba49 in DBA/p mice in, 875
 sarcoma T241 Lewis in C57BL/6JN mice in, 875
 calculation of average number of tumors per mouse in, 877-879
 counting and sizing of tumors in, 875
 dilution of tumor used in, 875
- Metastasis(es), spontaneous—(Contd.)
 limitation of metastases to lungs in, 875
 results for 5 experiments in, 876-881
 for amelanotic melanoma S91A in, 876
 for osteogenic sarcoma 112 in, 876, 877
 for primary S91 Cloudman melanoma in, 876, 877-878, 879
 for sarcoma dba49 in, 877, 878, 879, 880
 for sarcoma T241 Lewis in, 877, 878, 880
 to lymph nodes, and wound-washing results, after operation for cancer, lack of correlation between, 1042, 1043
 from cancer of vulva, 14-20
 in infiltrative squamous cell (epidermoid) vulvar carcinoma, significance of, 335
 to ovary, from advanced cancer of breast, and therapeutic oophorectomy, significance of, 126-130
- Methotrexate (4-amino-N¹⁰-methyl pteroylglutamic acid), continuous intra-arterial, antimetabolite-metabolite cancer chemotherapy using, with intermittent administration of intramuscular citrovorum factor (CF), report of, 744-752
 in treatment of malignant lymphomas in African children, use of, 266
 use of, in study of growth curves of human cancer after transplantation, report of, 1117-1121
- Methylcholanthrene, as carcinogen, development of carcinoma of gallbladder in dogs after implantation with, 1127-1130
 in experimental tumors of peripheral nerve trunk in rabbits, 1030-1035
 in study of progression of cervical dysplasia in mouse, 201-204
- 3-Methylcholanthrene (MC), as carcinogenic chemical, in newborn mice, report of, 308-317
- 2 α -Methyl dihydrotestosterone propionate hormonal therapy with, for metastatic carcinoma of female breast, 705-706
- MEYER, OVID O., *jt. su.*, 753-778
- Microscopy, interference, in study of exfoliated cells from experimental cancer of cervix, report of, 378-383
- MILLER, EDWARD, *jt. su.*, 744-752
- MILLER, THEODORE R., *jt. su.*, 1295-1300
- MILMORE, BENNO K., *jt. su.*, 622-628
- Mitomen (2,2-dichloro-N-methyl-N-oxide hydrochloride, Nitromin), in combined surgery and chemotherapy as treatment for malignant tumors, use of, 1197-1204
- MITRA, S., *jt. su.*, 502-506
- MODLIN, ROBERT K., and MORRIS, JOHN MCL.: The effect of certain chemical and physical agents on the radiation sensitivity of mouse tumors, 117-125
- MOERTEL, CHARLES G.; DOCKERTY, MALCOLM B., and BAGGENSTOSS, ARCHIE H.: Multiple primary malignant neoplasms. I. Introduction and presentation of data, 221-230
- DOCKERTY, MALCOLM B., and BAGGENSTOSS, ARCHIE H.: Multiple primary malignant neoplasms. II. Tumors of different tissues or organs, 231-237
- DOCKERTY, MALCOLM B., and BAGGENSTOSS, ARCHIE H.: Multiple primary malignant neoplasms. III. Tumors of multicentric origin, 238-248
- SAUER, WILLIAM G.; DOCKERTY, MALCOLM B., and BAGGENSTOSS, ARCHIE H.: Life history of the carcinoid tumor of the small intestine, 901-912
- MOFFITT, O. P., JR., *jt. su.*, 636-638
- MONELL, CHARLES M., *jt. su.*, 1131-1138
- MONSIE, BENITO, and WEINBERG, TOMAS: Cytochemical study of esterase activity of human neoplasms and stromal macrophages, 369-377
- MORAN, JOHN J.; BECKER, STANLEY M.; BRADY, LUTHER W., and RAMBO, V. BIRCH: Adenoid cystic carcinoma. A clinicopathological study, 1235-1250
- Morphology, of cells from ganglioneuroma of adrenal medulla, identical to hilus cells (extraparenchymal Leydig cells), report of, 421-425
- MORRIS, JOHN MCL., *jt. su.*, 117-125
- Mouse(mice), AKR, mature, failure to demonstrate acceleration of leukemia in, using cell-free brain filtrates from leukemic animals of same strain, report of, 496-501
- C3H, female, in study of progression of experimental cervical dysplasia, use of, 201-204
- C57BL/6JN, in study of spontaneous development of metastases after removal of "primary" tumor, use of, 875-882
- CDBA/F₁, in study of spontaneous development of metastases after removal of "primary" tumor, use of, 875-882
- combined action of cigarette tar and beta radiation on development of tumors in, 344-352
- DBA, tumors, effect of certain chemical physical agents on radiation sensitivity of, 117-125
- DBA/p, gray, in study of effect of growth hormone and adrenocorticotrophic hormone on transplanted tumors of, 507-511
- in study of spontaneous development of metastases after removal of "primary" tumor, use of, 875-882
- NH, white, in study of difference in host cell-virus relationship between tubular epithelium and stroma of kidneys infected with SE polyoma virus, 567-576
- Swiss, albino, in study of effects of carcinogenic chemicals, report on, 308-317
- (Millerton), in study of role of acidic fractions of tobacco in carcinogenesis, 1306-1315
- Mouth. See also Oral Cavity
- multiple carcinomas of, 512-516
- Mucin, in adenomatoid bronchial tumors, presence of, 172
- in carcinoma of pancreas, and multiple thrombi, relationship of, 950
- Mucosa(s), and submucosa, as sites of multiple lymphomatous polypoid of gastrointestinal tract, 247-256
- of bronchial tree, histological study of dysplastic changes in, 1226-1234
- of stomach and duodenum, metastases to, from carcinoma of breast, adrenal steroid-induced, report of, 426-431
- Multicentricity, of origin, of carcinoid tumors of small intestine, 902, 904
- MUIR, C. S.: Cancer in Singapore children, 534-538
- MURISON, P. J., *jt. su.*, 707-710
- Muscle(s), of cancer patients, excretion and distribution studies of radioactively-labeled alkylating drugs in, 955, 956
- MUSTACCHI, PIERO, and GRECO, TOMMASO: Angelo Nannoni on mastectomy (frontispiece), [4]
- MYERS, MAX H., *jt. su.*, 1041-1044
- Myleran, ¹⁴C-labeled, excretion and distribution studies in cancer patients with, 953-956
- ³⁵S-labeled, excretion and distribution studies in cancer patients with, 953-956
- Myxoma(s), in childhood, 629-635
 case reports, 631-633, 633-634
 clinical and pathological features in, 631
 gross characteristics of, 631
 criteria for diagnosis of, 629
 etiological factors in, 630-631
 anatomical distribution of tumors in, 630, 631
 patient age as, 630, 631
 race as, 630, 631
 sex as, 630, 631
 external soft tissues as location of, 633-634
 follow-up in, 630
 ileum as location of, 634
 jaws and maxillary sinus as locations of, 631-633
 microscopic appearance of, 631, 632, 633, 634
 orbit as location of, 634
 ovary as location of, 634
 results of treatment of, 634-635
 size of tumor in, 630
 odontogenic, in man, classification,

Myxoma(s), in childhood—(Contd.)
histopathology, and clinical behavior of, 89, 90

N-(2-naphthyl)-2-chloroacetamide (naphthylchloroacetamide), in chemotherapy of cancer, distribution of, 296-299

NADKARNI, MORRISWAR V.; **TRAMS, E. G.**, and **SMITH, PAUL K.**: Excretion and distribution studies with radioisotope-labeled alkylating drugs in cancer patients, 953-956

NAIB, ZUHER, M.: Exfoliative cytology of renal pelvic lesions, 1085-1087
Single trophoblastic cells as a source of error in the interpretation of routine vaginal smears, 1183-1185

NANNONI, ANGELO: On mastectomy, Frontispiece, [4]

Naphthol AS acetate, as substrate in cytochemical study of esterase activity in human neoplasms and stromal macrophages, report of, 369-377

Naphthylchloroacetamide (*N*-(2-naphthyl)-2-chloroacetamide), in chemotherapy of cancer, distribution of, 296-299

activity in epithelial cells in, 298
determination of enzymatic activity in, 296-297

human liver as most active tissue in, 297, 298

hydrolysis by human organs and serum in, 297-298

mouse tissue hydrolysis of, 297

preparation of tissue homogenate in, 296

types of disease and activity in, 297

NASH, ARTEMIS, and **STOUT, ARTHUR PURDY**: Malignant mesenchymomas in children, 524-533

NASIELL, MAGNUS: Histiocytes and histiocytic reaction in vaginal cytology. A survey of the risks of false positive diagnosis, 1223-1225

Neoplasm(s). *See also specific names and sites*
human, cytochemical study of esterase activity of, 369-377

Nerve(s), trunks of, peripheral, in rabbits, experimental tumors of, 1030-1035

development of palpable tumors in, 1030-1035

methylcholanthrene, in acetone, used as carcinogenic agent in, 1030

dosage used in, 1030

pathological findings in, 1033-1035

comparison of preliminary study in rats and, 1033, 1035

gross examination in, 1031, 1033

involvement of nerve roots within spinal canal in, 1031, 1033

histological appearance of thickened nerve trunks in, 1033

microscopic findings in, 1031-1035

areas suggestive of dermatofibrosarcoma protuberans and areas of undifferentiated sarcomas in, 1032, 1033

cells similar to Vater-Pacini's corpuscles in, 1032, 1033

malignant or benign appearance of tumors in, 1031, 1032, 1033-1035

pattern of nuclei similar to that of schwannomas of Antoni type A in, 1031, 1033, 1035

sheets of cells similar to those in epithelial tumors or meningiomas in, 1032, 1033

surgical exposure of sciatic nerves in, 1030

Neurosurgery, in treatment of adenoid cystic carcinoma, report of, 1248

Nevus(i), plantar and palmar, prophylactic removal of, dissenting view on, 102-104

alternative approach to management of, 103

clinical observation as factor in, 103

morbidity rate involved in removal of plantar and palmar nevi as factor in, 103

over-all incidence of nevi in group studied in, 103

sex of individuals in, 103

Nipple(s), Paget's disease of, some observations on, 653-672

growth by ascent or descent in, 653-672

Nitrogen mustard (HN2), in therapy of giant follicular lymphoblastoma, report of, 771, 772-773

Nitroimin (2,2-dichloro-*N*-methyl-*N*-oxide hydrochloride, Mitomen), in combined surgery and chemotherapy as treatment for malignant tumors, 1197-1204

Nose, as location of congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, progionoma, and pigmented epulis of infancy, 1090-1108

Nucleus(i), and cells, diameters of, in study of measurement of quantitative characteristics of exfoliated cells from female genital tract, report of, 437-453

Nutrition, as cause of cirrhosis, autoradiographic and histological study of repair after partial hepatectomy, report of, 963-967

effect of, on hepatic metastases, experimental studies of, 547-554

Occupation(s), as etiological factor in cancer of esophagus, study of, 395, 397, 398, 401

O'Connor, G. T.: Malignant lymphoma in African children. II. A pathological entity, 270-283

jt. sw., 258-269

Odontoma(s), ameloblastic, in man, classification, histopathology, and clinical behavior of, 87-88

complex, in man, classification, histopathology, and clinical behavior of, 88-89

compound, in man, classification, histopathology, and clinical behavior of, 89-90

in man, general classification, histopathology, and clinical behavior of, 87

OGURA, JOSEPH H., *jt. sw.*, 55-66

Oophorectomy, therapeutic, for advanced cancer of breast, significance of metastases to ovary and of ovarian cortical stromal hyperplasia in, 126-130

Oral cavity. *See also specific sites of organs*

as location of congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, progionoma, and pigmented epulis of infancy, report of, 1090-1108

carcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 238-239

exfoliated normal and malignant cells, effects of radiation on, 587-596

Orbit, as location of myxoma in children, report of, 634

Organ(s). *See also specific names of organs*

endocrine, variation in weight of, in laboratory animals, and leukemia incidence curve, possible relationship of, in epidemiological study of seasonal incidence of acute leukemia, 1303, 1305

OSBORNE, MELVIN P., and **PITTS, R. MARSHALL**: Therapeutic oophorectomy for advanced breast cancer. The significance of metastases to the ovary and of ovarian cortical stromal hyperplasia, 126-130

Osteitis deformans, complicated by superimposed bone sarcomas, 517-523

fibrosis, in acute leukemia, 614

Osteoporosis, in acute leukemia, report of, 613-614

OTIS, RICHARD D., *jt. sw.*, 913-920

Ovary(ies), adenocarcinomas of, treated by 5-fluorouracil (5-FU), clinical evaluation of, 138, 139, 140-145, 150-151

and uterus, adenocarcinomas of, occurring as coexistent or sequential primary neoplasms, report of, 817-826

case reports, 818-823

criteria used in classification in, 823-824

microscopic appearance of tumors in, 818-823

operative procedures used in, 818-823

pathological findings in, 818-823

possible factor of hormonal disturbance in, 824

preparation of tissue sections in, 818

principal characteristics of, 818-825

Ovary(ies), adenocarcinoma of—(Contd.)
clinical findings in, 817, 818-823

prognosis in, 818-825
as location of myxoma in children, report of, 634

bilateral carcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 243-244

cancer of, developing in dermoid cysts of, 989-1000

background data for, 989-990
age of patients with benign dermoid cysts in, 989

gross changes suggesting malignant change in, 989
structure of dermoid cysts in relation to types of cancer that may develop in, 989

symptoms of benign dermoid cysts in, 989
case reports, 990-994

extent of tumor at time of operation in, 990-994
pathological findings in, 991-999

prognosis in, 991-999
size of tumor in, 990-994

symptoms and signs in, 990-994
treatment in, 989-994

type of cancer in, 990-999
carcinoid in, 990, 993-994, 995-999

combination of squamous cell carcinoma and trabecular adenocarcinoma in, 990, 995-999

leiomyosarcoma in, 990, 992-993, 995-999

spindle cell sarcoma in, 990, 995-999

squamous cell type of carcinoma in, 990-992, 995-999
undifferentiated carcinoma type in, 990, 995-999

metastatic, luteinization of stroma of, endocrine significance of, 827-840

case reports, 835-836
categories of ovarian metastatic disease included in, 827

correlation of pathological and endocrine findings in, 833-835

abnormal vaginal bleeding in, 833
appearance of endometrium in, 833

association of increased libido with, 833
of virilization with, 833, 837-839

operative procedures used in, 835-836
pathological findings in, 827-831

by location of metastases in, 827-831, 831-833
comparison with "foam" cells in, 831

nature of lutein-like cells in, 831-833
role of metabolism of steroid hormones in, 837-839

similarity between cells found in cases of hyperthecosis and, 837-839

symptoms in, 835-836
malignant lymphomas of, in African children, pathological entity of, 275, 276

pelvic and abdominal carcinomatosis primary in, attempted palliation by radical surgery for, 384-388

"carcinomatosis ileus" in, 384
case reports, 385, 386-387

excision of markedly involved omentum with or without resection of peritoneal and mesenteric implants in, 385

intestinal obstruction in, 384
previous therapy received in, 384

resection of involved segment of sigmoid or descending or transverse colon, with or without resection of omentum, with or without resection of segment of small bowel, and with or without excision of larger peritoneal implants in, 385

of omentum heavily infiltrated by cancer and radical pan-hysterectomy in, 385

- Ovary(ies) pelvic and abdominal, carcinomatosis primary in—(Contd.)
 of one or more segments of small bowel, with or without omentectomy and excision of larger peritoneal and mesenteric implants in, 385
 right colectomy, with or without omentectomy and with or without resection of larger peritoneal metastases or other segments of bowel in, 385-386
 surgical mortality rate in, 387
 survival of patients in, 385-388
 temporary amelioration of symptoms in, 384
- OZELLO, LUCIANO, and STOUT, ARTHUR PURDY: The epithelial origin of giant cell carcinoma of the lung confirmed by tissue culture. Report of a case, 1052-1056
- PACK, GEORGE T., and MILLER, THEODORE R.: Middle hepatic lobectomy for cancer, 1295-1300
- Paget's disease, complicated by bone sarcoma, report of, 517-523
 case reports, 517-520
 prognosis in, 523
 survival time in, 517-523
 of nipple, observations on, 653-672
 ascent or descent within ducts in, 653, 655
 aspects of architecture of mammary duct system in, 657-659
 basal zone in, 657-659
 eosinophilic staining in, 657-659
 mammary epithelium in, 657-659
 concentration of Paget cells at epidermal-dermal junction in, 655
 controversy over second invasive growth in, 653
 extension of Paget cells in, 657
 findings for Paget cells in, 654, 655-672
 bizarre multinucleated forms in, 654, 655
 degree of staining of cells in, 654, 655, 656
 hyperchromasia in, 654, 655
 melanin granules in, 654, 655
 mitotic figures in, 654, 655
 nuclei in, 654, 655
 shrunken central nucleocytoplasmic mass in, 654, 655
 size of cells in, 655
 measurement of primary carcinoma in, 655
 nuclei in, 655
 relationship between deeply situated cancer and Paget infiltration within epidermis in, 661-665
 avenue of extension in, 661
 coincidental finding of metaplasia within tributary duct in, 665
 possibility of diffuse intraductal neoplastic transformation rather than metastatic spread in, 665
 structure of lactiferous ducts and, 661-665
 relationship between lesion within depths and that within epidermis in, 655
 serial sectioning of nipple in, 655
 spread by permeation in, 655-672
 underlying carcinoma in, 659-661
 nuclear and cytoplasmic structure in, 659-661
 tendency to retraction in, 661
 of vulva, differentiation between Bowen's type of squamous cell carcinoma and, 318, 329
- Pain, bone, in acute leukemia, skeletal lesions and, 608-621
- Palate, hard, as site of origin of adenoid cystic carcinoma, 1235-1250
- Palliation, by radical surgery, for pelvic and abdominal carcinomatosis primary in ovaries, report of, 384-388
 of metastatic breast carcinoma, alterations in serum protein levels of patients receiving, 1081-1084
- Pancreas, carcinomas of, and multiple thrombi, morphological study of, 944-952
 classification of cases in, 944, 947
- Pancreas, carcinomas of—(Contd.)
 control series in, 947
 microscopic findings in, 947-951
 "differentiating ductal carcinomas" in, 947
 "pure ductal carcinomas" in, 947
 "well differentiated islet cell carcinomas" in, 948
 mucin in relation to incidence of multiple thrombi in, 950
 multiple thrombi in relation to histological type and location of, 948
 relationship of carcinomatous metastases to development of multiple thrombi in, 948-949
 site of tumor in relation to incidence of multiple thrombi in, 949-950
 as multiple primary malignant tumors of multicentric origin, report of, 244
 involvement of, by giant follicular lymphoblastoma, report of, 762
 malignant lymphomas of, in African children, pathological entity of, 275
- Panhysterectomy, simple total, inadequacy of, in management of invasive cervical carcinoma, report of, 582-586
- Papilloma(s), laryngeal, treatment of, with bovine wart vaccine, 636-638
- Paraganglioma(s), nonchromaffin, comparative study of, 779-787
 clinical observations in, 780, 782
 for carotid body tumors in, 780, 782
 for jugular body tumors in, 782
 symptoms in, 780, 782
 pathological findings in, 782-785
 gross description of, 782
 alveolar soft part sarcoma in, 784
 type of rhabdomyosarcoma in, 784-785
 hemangioblastoma of cerebellum in, 784
 microscopic description of, 782-785
 aortic body tumor in, 784
 periodic acid-Schiff (PAS) reaction in, 784-787
 retroperitoneal paraganglioma in, 784, 786
 special stains and histochemical techniques used in, 784-787
 possible histogenesis in, 785
 staining techniques used in, 780
 retroperitoneal, report of, 784, 786
- Paraplegia, resulting from malignant lymphomas, in African children, clinical syndrome of, 264-265
- Parasite(s), in relation to incidence of malignant lymphoma in African children, possible "priming" action of, 282
- Parathyroid, function of, in advanced cancer, hypercalcemia and, relationship of, between, 934-943
- PARKER, JAMES E., *jt. au.*, 213-216
- Parotid gland, involvement of, in giant follicular lymphoblastoma, 758
- PARSONS, LANGDON, *jt. au.*, 42-54
- PECHET, GISELLE, and MACDONALD, RICHARD A.: Repair of nutritional cirrhosis. Autoradiographic and histological study after partial hepatectomy, 963-970
- Pedigrees, in studies of inheritance of cancer of human uterine cervix, use of, 180, 181
- Pelvis, carcinomatosis of, primary in ovaries, attempted palliation by radical surgery in, 384-388
 renal, lesions of, exfoliative cytology in, 1085-1087
 collection of urine in, 1086
 correlation of cytological findings with surgical findings in, 1086
 cytological interpretation of upper urinary tract lesions in, 1086
 possible factors affecting interpretation of cells in, 1086
 preparation of patient in, 1085-1086
 staining and mounting of filter in, 1086
 types of lesions studied in, 1086
 use of Millipore filter apparatus in, 1086
 value, as additional diagnostic tool, of, 1087
- Pelvis, carcinomatosis of—(Contd.)
 wide variation in published results of cytological examinations of urine in, 1085
- Pericardium, involvement of, by giant follicular lymphoblastoma, 759
- PERLIA, CHARLES P.; KOFMAN, SYDNEY; SKY-PECK, HOWARD, and TAYLOR, SAMUEL G., III: Clinical use of 6-aminonicotinamide in patients with disseminated neoplastic disease, 644-648
- Pharynx, epitheliomas of, as multiple primary malignant tumors of multicentric origin, report of, 239
- PHI. See Phosphohexose isomerase
- Phosphate(s), renal, excretion of, in advanced cancer, effect of citrate infusion on, 934-943
- Phosphohexose isomerase (PHI), levels of, in serum of patients with carcinoma of breast, report of, 726-733
- PICKREN, JOHN W.: Significance of occult metastases. A study of breast cancer, 1266-1271
- PIERCE, G. B., JR., and VERNER, E. L.: An in vitro and in vivo study of differentiation in teratocarcinomas, 1017-1029
- PIERCE, MILA I., *jt. au.*, 496-501
- PIETRA, GIUSEPPE; RAFFAPORT, HENRY, and SHUBIK, PHILIPPE: The effects of carcinogenic chemicals in newborn mice, 308-317
- PINDBOG, JENS J., *jt. au.*, 73-101
- PINKEL, DONALD: Chlorambucil in childhood cancer, 36-41
- PITTS, R. MARSHALL, *jt. au.*, 126-130
- Plummer-Vinson disease, as etiological factor in cancer of esophagus, study of, 391, 402-403, 407, 410
- Pneumonectomy, for primary Hodgkin's disease of lung, results of, 1151-1165
- Pneumonitis, postoperative, use of pancreatic deoxyribonuclease (dornase) in treatment of, 415-416, 420
- Polyoma virus, infections by, relationship of, to leukemia in mice, 496-501
- Polypoid, multiple, lymphomatous, of gastrointestinal tract, report of, 249-257
- Potaba. See Potassium β -aminobenzoate
- Potassium β -aminobenzoate, prolonged remission of reticulum cell sarcoma complicating atopic dermatitis, after therapy with, 5-12
- Predetermination, biological, and breast cancer, re-evaluation of, 1122-1126
- PRESSMAN, JOEL J.; SIMON, MILDRED; BURKE, and MONELL, CHARLES M.: Anatomic studies related to the dissemination of cancer of the larynx, 1131-1138
- PRIMIKIRIOS, NICHOLAS, *jt. au.*, 597-607
- Prognosis(es), of gelatinous carcinoma of breast, report of, 699-704
- Progonoma(s), congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, and pigmented epulis of infancy, report of, 1090-1108
 melanotic, in scapula region, report of, 1088-1089
 microscopic findings in, 1088-1089
- Prostate, male, cancer of, development of breast cancer after estrogen treatment for, 234
- Protein, content of, in diet, effect of on hepatic metastases, studies of 547-554
- PRZYBORA, LUCIAN A.: Leiomyosarcoma in situ of the uterus, 483-492
- Pseudosarcomatous fasciitis, in children, report of, 1216-1222
 age and sex of patient and size of tumor in, 1216-1217, 1219, 1221-1222
 case reports, 1221-1222
 clinical data in, 1219
 diagnoses made in, 1217
 location of tumor in, 1217, 1219, 1221-1222
 microscopic features in, 1217-1222
 foci of calcification seen in, 1219, 1220, 1221
 invasion of surrounding tissue in, 1218, 1219
 pattern of tumor behavior in, 1217
 symptoms in, 1217, 1219, 1221-1222
 treatment of choice in, 1219, 1221-1222
- Psychology, in management of terminal cancer patients, role of, 1063-1073
 some observations about patients with

- Psychology—(Cont'd.)
breast mass based on, 979-988
Public education, about breast cancer, effect of, on patients with breast mass, report on, 985
relation of delay in seeking treatment and, 985
- Rabbit(s), peripheral nerve trunks of, experimental tumors of, 1030-1035
Radiation. See also specific forms of radiation
as therapy for cancer of vulva, 13-20
survival rates after, 15-20
contraindications for, 17-18
for malignant lymphomas in African children, use of, 266
as treatment for multiple oral carcinoma, use of, 515
beta, and cigarette tar, in mice, combined action of, 344-352
deep, as therapy for multiple lymphomatous polyposis of gastrointestinal tract, report of, 256
effects of, on exfoliated normal and malignant oral cells, 587-596
in relation to therapy with 5-fluorouracil (5-FU) for cancer, clinical evaluation of, 131-152
in treatment of cervical cancer, development of in situ epidermoid carcinoma after, 353-360
of nonbenign bronchial carcinoids and cylindromas, 544-546
ionizing, therapeutic, for initial malignant tumor, induction of second primary cancer by, 232-233
therapy, and later development of Bowen's type of intraepithelial and infiltrative carcinoma of vulva, possible relationship of, 321-328
for management of intracranial metastases, value of, 577-581
inadequacy of, in management of invasive cervical carcinoma, report of, 582-586
Radiation response (RR), of canine small intestine, effect of succinylsulfathiazole (Sulfasuxidine) on, 862-874
of exfoliated normal and malignant oral cells, report of, 587-596
Radioactive labeling, of alkylating drugs, excretion and distribution studies of cancer patients using, 953-956
strontium (^{90}Sr), use of, for beta radiation in mice, combined action of cigarette tar and, 344-352
Radiosensitivity, of mouse tumors, effect of certain chemical and physical agents on, 117-125
Radiotherapy, for adenoid cystic carcinoma, report of, 1247-1248
RAMBO, V. BLUCH, *jt. au.*, 1235-1250
RAPAPORT, HENRY, *jt. au.*, 308-317
Rat(s), Lewis strain, adult, cancericidal effects of X irradiation at body temperature below 10°C , on Lewis lymphoma in, 1060-1062
Sprague-Dawley, female albino, in study of effects of "aging" on development of tumors after inoculation of carcinosarcoma 256 Walker cells, report of, 1057-1059
male, in autoradiographic and histological study of repair of nutritional cirrhosis after partial hepatectomy, use of, 963-970
use of, in experimental study of effect of nutrition on hepatic metastases, report of, 547-554
Wistar, adult, female, in study of effect of human fibrinolysin on survival of carcinosarcoma 256 Walker cells in blood, 957-962
RATZKOWSKI, E., and HOCHMAN, A.: Survival of patients with "recurrent" or inoperable carcinoma of the breast with special consideration of the effect of hormonal treatment. A study of 338 cases, 300-307
RAZIS, DENNIS V.; AMBRUS, JULIAN L.; ROSS, CHARLES A.; STUTZMAN, LEON; SOKAL, JOSEPH E.; REJALI, ABDES M., and BARROS, THOMAS T.: Use of AB-103 (a "dual antagonist") in the treatment of bronchogenic carcinoma, 853-861
Rectum. See also Anus and Colon adenocarcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 240
Recurrence, of cervical carcinoma, after simple total hysterectomy, management of, 582-586
of mesenchymomas, in children, incidence of, 526-533
Reductase, glutathione (GR), levels of, in serum of patients with carcinoma of breast, report of, 726-733
REID, J. D., and CARR, A. H.: The validity and value of histological and cytological classifications of lung cancer, 673-698
REJALI, ABDES M., *jt. au.*, 853-861
Religion, as etiological factor in cancer of esophagus, 393, 398-400, 404, 409
Respiratory tract. See specific names of organs
cancer of, etiological factors in, 410
Reticuloendothelial system. See also specific names of organs
malignant lesions of, as multiple primary malignant lesions of multicentric origin, report of, 244-245
Rhabdomyosarcoma(s), alveolar type, report of, 784-785
in children, as part of malignant compound tumor, 524-533
RICHARDSON, GEORGE S., *jt. au.*, 827-840
RIGGENBACH, NEIL, and VON KAULLA, KURT, N.: Urokinase excretion in patients with carcinoma, 889-896
ROBERTS, GUY F., *jt. au.*, 699-704
ROBERT, J.; VAN RYMENANT, M., and LAGAE, F.: Enzymes in cancer. III. Triosephosphate isomerase activity of human blood serum in normal individuals and in individuals with various pathological conditions, 1166-1174
ROMSDAHL, MARVIN D.; CHU, ELIZABETH W.; HUME, ROXANE, and SMITH, ROBERT R.: The time of metastasis and release of circulating tumor cells as determined in an experimental system, 883-888
RONGONE, E. L., *jt. au.*, 195-198, 707-710
ROPER, CHARLES L., *jt. au.*, 1251-1258
ROSE, ARTHUR; WEST, MICHAEL, and ZIMMERMAN, HYMAN J.: Serum enzymes in disease. V. Isocitric dehydrogenase, malic dehydrogenase, and glycolytic enzymes in patients with carcinoma of the breast, 726-733
ROSS, CHARLES A., *jt. au.*, 853-861
ROSS, NEAL L., *jt. au.*, 13-20
ROSVOLL, RANDI V., *jt. au.*, 734-743
ROTHENBERG, ALBERT: Psychological problems in terminal cancer management, 1063-1073
ROTKIN, I. D.: Studies on the inheritance of cancer of the human uterine cervix, 179-194
RR. See Radiation response
RUBENSTON, ALBERT I., *jt. au.*, 21-29, 30-35
RUBIN, CYRUS E., *jt. au.*, 1074-1080
Salivary gland(s), major, as site of origin of adenoid cystic carcinoma, report of, 1235-1250
malignant lymphomas of, in African children, clinical syndrome of, 265
pathological entity of, 275
type of tumors, bronchial, 161-178
Saltzstein, Sidney L., *jt. au.*, 862-874
Sanamycin (actinomycin C), in combined surgery and chemotherapy for malignant tumors, 1197-1204
SAPHIR, OTTO, *jt. au.*, 21-29, 30-35
Sarcoma(s). See also specific names and sites
alveolar soft part, report of, 782, 784, 785-787
ameloblastic, in man, classification, histopathology, and clinical behavior of, 85-86
bone, complicating Paget's disease, report of, 517-523
case reports, 517-520
survival time in, 517-523
db49, in experimental study of effectiveness of small amounts of preoperative irradiation in preventing growth of tumor cells disseminated at surgery, report on, 284-295
in spontaneous development of Sarcoma(s), db49—(Cont'd.)
metastases after removal of "primary" tumor, use of, 875-882
transplanted in mice, effect of growth hormone and adrenocorticotrophic hormone on, 507-511
in situ, development of, in patients with uterine myomas, 483-492
osteogenic, in children, as part of malignant compound tumor, report of, 524-533
112, in study of spontaneous development of metastases after removal of "primary" tumor, use of, 875-882
reticulum cell, complicating atopic dermatitis, prolonged remission of, after administration potassium β -aminobenzoate (KPAW, Potaba) therapy, report of, 5-12
case report, 5-11
diagnosis in, 5, 6, 9
dosage used in, 9
histological findings in, 6-9
other conditions exhibiting benefit after use of, 5, 11
5-year survival in, 9-11
in children, as part of malignant compound tumor, 524-533
presenting as multiple polyposis of gastrointestinal tract, report of, 249-257
spindle cell, developing in dermoid cysts of ovary, 990, 995-999
synovial, in children, as part of malignant compound tumor, 524-533
T241 Lewis, in spontaneous development of metastases after removal of "primary" tumor, 875-882
use in determining time of metastasis and release of circulating tumor cells as determined in experimental study, 883-888
use in experimental study of effectiveness of small amounts of preoperative irradiation in preventing growth of tumor cells disseminated at surgery, report on, 284-295
undifferentiated, in children as part of malignant compound tumor, report of, 524-533
SAUER, WILLIAM G., *jt. au.*, 901-912
Scapula, melanotic progionoma in region of, 1088-1089
case report, 1088
microscopic findings in, 1088-1089
columnar epithelium in, 1088, 1089
lack of "retinal anlage" or "ameloblastic" structures in, 1088
Scar(s), peripheral lung cancer arising in, report of, 1205-1215
SCHATZKI, STEFAN C., and DUDLEY, H. ROBERT: Bone sarcoma complicating Paget's disease. A report of 3 cases with long survival, 517-523
SCHULMAN, HAROLD, and CAVANAGH, DENIS: Intraepithelial carcinoma of the cervix. The predictability of residual carcinoma in the uterus from microscopic study of the margins of the cone biopsy specimen, 795-800
SCHWARTZ, MORDECAI: A biostatistical approach to clinical tumor growth, 1272-1294
Scintiscanning, in preoperative selectivity of patients with nodular thyroids, use of, 68-69
scintiscanning alone as, 68-69
use of radioactive iodine (^{131}I) in, 68, 69
SCULLY, ROBERT E.: Spermatocytic seminoma of the testis. A report of 3 cases and review of the literature, 788-794
and COHEN, RICHARD B.: Ganglioglioma of adrenal medulla containing cells morphologically identical to hilus cells (extraparenchymal Leydig cells), 421-425
and RICHARDSON, GEORGE S.: Luteinization of the stroma of metastatic cancer involving the ovary and its endocrine significance, 827-840
jt. au., 989-1000
SEGALOFF, ALBERT; WEETH, J. B.; RONGONE, E. L., and MURISON, P. J.: Hormonal therapy in cancer of the breast. XVIII. Effect of androstenediol dipropionate (5a-an-

- drostan-3 α ,17 β -diol dipropionate) therapy on clinical course and hormonal excretion, 707-710
- WERTH, J. B.; RONGONE, E. L., and BOWERS, C. Y.: Hormonal therapy in cancer of the breast. XVII. The effect of androsterone on clinical course and hormonal excretion, 195-198
- Segmentectomy(ies), median left, for liver cancer, report of, 1295-1300
- Selectivity, preoperative, in patients with nodular thyroids, criteria employed in, 68-71
- SELIGMAN, ARNOLD M., *jt. au.*, 296-299
- Seminoma(s), canine, similarity of, to classical and spermatocytic seminoma, report of, 793-794
- classical, compared to spermatocytic seminoma of testis, 788-794
- spermatocytic, of testis, 788-794
- Serial step sectioning, after cervical biopsy with abnormal findings, practicality of, 213, 215-216
- Serotonin, in adenomatoid bronchial tumors, role of, 172, 173
- production of, in functioning carcinoid tumors of small intestine, report of, 908
- Serum(s), blood, human, of normal individuals and individuals with various pathological conditions, triosephosphate isomerase activity of, 1166-1174
- description of enzyme in, 1166
- effect of hemolysis in, 1168
- of ions in, 1168
- of pH in, 1168
- of temperature in, 1168-1169
- procedure followed in, 1167
- reagents employed in, 1166-1167
- composition of reaction mixture in, 1166-1167
- preparation of buffer in, 1167
- results for normal individuals in, 1167, 1169, 1174
- variation in values found in, 1167, 1169, 1174
- for patients with various diseases in, 1168, 1169-1174
- classification of patient groups in, 1169
- generalized cancer as, 1169
- localized cancer as, 1169
- miscellaneous conditions as, 1169
- non-neoplastic liver injuries as, 1169
- levels for patients with cirrhosis, obstructive jaundice, and/or hepatitis in, 1172, 1173, 1174
- with generalized cancer and no liver metastases in, 1171
- with generalized cancer and possible liver metastases in, 1170-1171
- with generalized cancer and probable liver metastases in, 1169-1170
- with leukemias in, 1171-1172
- with localized cancer in, 1172
- with miscellaneous conditions in, 1172, 1173
- proof of liver metastases in, 1169
- unit of activity measurement employed in, 1166
- equation used in, 1166
- of patients with carcinoma of breast, levels of enzymes in, 726-733
- proteins, in patients receiving palliative therapy for metastatic breast carcinoma, alterations in, 1081-1084
- SHARP, GEORGE S.; BULLOCK, WELDON K., and HELSPER, JAMES T.: Multiple oral carcinomas, 512-516
- SHEEHAN, F. R., *jt. au.*, 205-209
- SHELINE, GLENN E., *jt. au.*, 587-596
- SHERLOCK, PAUL, *jt. au.*, 1117-1121
- SHERMAN, JOSEPH D., *jt. au.*, 1117-1121
- SHOLTON, LEON J.; INCZE, JOSEPH S., and WERK, EMILE E., JR.: Adrenocortical width in carcinoma of the lung, 105-110
- SHUBIK, PHILIPPE, *jt. au.*, 308-317
- SILVERMAN, SOL, and SHELINE, GLENN E.: Effects of radiation on exfoliated normal and malignant oral cells. A preliminary study, 587-596
- SIMON, MILDRED BURTZ, *jt. au.*, 1131-1138
- Sinus(es), maxillary, as location of myxoma in children, report of, 633-634
- Skeleton. *See also specific anatomical sites*
- lesions, in acute leukemia, 608-621
- malignant lymphomas of, in African children, clinical syndrome of, 265
- pathological entity of, 272
- Skin, basal cell carcinoma of, metastasizing, report of, 1036-1040
- incidence of metastases in, 1036-1040
- location of metastases in, 1036-1040
- time of metastases in, 1036-1040
- delayed test responses of, in cases of Hodgkin's disease and lymphosarcoma, report of, 597-607
- epitheliomas of, as multiple primary primary malignant tumors of multicentric origin, report of, 238
- involvement of, in giant follicular lymphoblastoma, 756-757
- tumors of, development of, in mice, after exposure to cigarette tar and beta radiation, 344-352
- Skull, as location of congenital melanocarcinoma, melanotic adamantinoma, retinal anlage tumor, progonoma, and pigmented epulis of infancy, report of, 1090-1108
- SKY-PECK, HOWARD, *jt. au.*, 644-648
- Smear(s), cervical, in study of suitability of quantitative cytological measurements for automatic prescreening of exfoliated cells from female genital tract, report of, 455-468
- vaginal, routine, single trophoblastic cells as source of error in interpretation of, 1183-1185
- SMITH, PAUL K., *jt. au.*, 953-956
- SMITH, ROBERT R., *jt. au.*, 284-295, 883-888, 1041-1044
- Smoking. *See also Cigarette(s) and Tobacco*
- acidic fractions involved in, as promoters in carcinogenesis, role of, 1306-1315
- as etiological factor in cancer of esophagus, study of, 389-413
- effect of on basement membrane and elastic fibers of trachea and bronchus, report of, 1175-1182
- histological study of dysplastic changes in mucosa of bronchial tree and, 1226-1234
- Sodium fluoride, inhibitory effect of, on esterase activity of human neoplasms and stromal macrophages, report of, 373, 375
- taurocholate, inhibitory effect of, on esterase activity of human neoplasms and stromal macrophages, report of, 373, 375
- SOKAL, JOSEPH E., and PRIMIKIRIOS, NICHOLAS: The delayed skin test response in Hodgkin's disease and lymphosarcoma. Effect of disease activity, 597-607
- jt. au.*, 853-861
- SOLARIC, S., *jt. au.*, 344-352
- SOMMERS, SHELDON C., *jt. au.*, 1117-1121
- SOUTHAM, CHESTER, M., and BRUNSCHWIG, ALEXANDER: Quantitative studies of autotransplantation of human cancer. Preliminary report, 971-978
- SPJUT, HARLAN J.; ROPER, CHARLES L., and BUTCHER, HARVEY R., JR.: Pulmonary cancer and its prognosis. A study of the relationship of certain factors to survival of patients treated by pulmonary resection, 1251-1258
- Spleen, involvement of, by giant follicular lymphoblastoma, 761-762
- malignant lymphomas of, in African children, pathological entity of, 275
- SPRATT, JOHN S., JR.; HEINBECKER, PETER, and SALZSTEIN, SIDNEY L.: The influence of succinylsulfathiazole (Sulfasuxidine) upon the response of canine small intestine to irradiation, 862-874
- SP⁹⁰. *See* Radioactive strontium
- STABENAU, JAMES R., *jt. au.*, 608-621
- STAHL, THEODORE, *jt. au.*, 67-72
- Staining, of carcinoids of bronchus, reactions in, 171-173
- argyrophil-positive cells in, 171-173
- argyrophil-positive cells in, 171-173
- use of silver techniques in, 171
- Statistics. *See also Epidemiology*
- analysis of 5 "stomach cancer families" in State of Utah, 1005-1016
- biometric approach to etiology of cancer of cervix, 153-161
- declining rate of increase in mortality attributed to carcinoma of lung, report of, 622-628
- epidemiological study of seasonal incidence of acute leukemia, 1301-1305
- etiological factors in cancer of esophagus, study of, 389-413
- for breast cancer, biological preterminism and, re-evaluation of, 1122-1126
- hereditary influences in patients with multiple primary malignant tumors, 234-235
- incidence of cancer in Singapore children, report of, 534-537
- of clinical syndrome of malignant lymphoma in African children, report of, 258-269
- of multiple primary malignant tumors of different tissues or organs, report of, 231-237
- of multicentric origin, 238-248
- over-all report of, 221-230
- of pathological entity of malignant lymphoma in African children, report of, 270-283
- of stomach cancer in spouses of patients with, 199-200
- studies on inheritance of cancer of human uterine cervix, 179-194
- STERN, ELIZABETH, and DIXON, W. J.: Cancer of the cervix—a biometric approach to etiology, 152-160
- Steroid(s). *See also Hormone(s) and specific names*
- adrenal, role of, in development of gastroduodenal metastases from carcinoma of breast, 426-431
- STEWART, SARAH E., *jt. au.*, 567-576
- Stewart-Treves syndrome, role of, in incidence of multiple primary malignant tumors, report of, 233-234
- Stomach, adenocarcinomas of, as multiple primary malignant tumors of multicentric origin, 239
- treated by 5-fluorouracil (5-FU), evaluation of, 138, 140-145
- cancer, in families in State of Utah, analysis of, 1005-1016
- ascertainment of kindreds in, 1007, 1009
- selection of 5 proband families to be studied in, 1007, 1009
- definition of family in, 1009
- of kindred in, 1009
- determination of medical histories for kindred members in, 1009
- diminution trend noted in each kindred in, 1015
- results for 5 proband families in, 1010-1013
- significance of high incidence of stomach cancer found in, 1013, 1015
- in spouses of patients with, incidence of, 199-200
- etiological exogenous factors in home environment, 199
- frequency of stomach cancer deaths and other cancer deaths in spouses of stomach cancer patients and controls in, 200
- new method of selecting controls in, 199-200
- types of cancer deaths occurring in spouses of stomach cancer patients and controls in, 199
- use of death certificate information in, 199-200
- simplified chymotrypsin technique for exfoliative cytology in, 1074-1080
- accuracy of diagnosis by 1077-1078
- acetate buffer as lavage fluid in, 1074-1075
- comparison of results by "long" and "short" methods in, 1077-1078

Stomach, cancer—(Contd.)

- diagnostic accuracy of chymotrypsin lavage in, 1075
influence of collection methods on case of cancer diagnosis in, 1078
"blind" evaluation of, 1078
main deterrents to wide application of gastric diagnosis and, 1074
original chymotrypsin method used in, 1076-1077
aspiration of fluid in, 1076
centrifugation of fluid in, 1076
fixation of slides in, 1076
injection of Ringer's solution into patient's stomach in, 1076
instillation of crystalline alpha chymotrypsin in, 1076
number of slides made in, 1076-1077
preparation of smears in, 1076
other methods used in diagnosis of, 1079-1080
preparation of patient's stomach in, 1076
importance of hydration in, 1076
previous lavage techniques tried in, 1074
proof of benign or malignant nature of gastric lesion in, 1075
short chymotrypsin method used in, 1077
cell collection in, 1077
preparation of slides in, 1077
use of chilled Levin tube in, 1076
of crystallized alpha chymotrypsin in, 1074
metastases to, from carcinoma of breast, adrenal steroid-induced, report of, 426-431
STOUT, ARTHUR PURDY: Fred W. Stewart, M.D., Ph.D.: Editor of Cancer, Frontispiece, [1148-1149]
Pseudosarcomatous fasciitis in children, 1216-1222
jt. au., 469-482, 524-533, 629-635, 1052-1056, 1186-1196
Stroma, cervicovaginal, radiation changes in, after radiotherapy for cervical cancer, report of, 359
in squamous cell carcinoma of esophagus, reaction of, prognostic factor, report of, 921-933
luteinization of, in metastatic ovarian cancer, endocrine significance of, 827-840
of mouse kidney, infected with SE polyoma virus, and tubular epithelium, difference in host cell-virus relationship in, report of, 567-576
ovarian cortical, hyperplasia of, from advanced cancer of breast in relation to therapeutic oophorectomy, significance of, 126-130
STUTZMAN, LEON, *jt. au.*, 853-861
Submucosa, and mucosa, as sites of multiple lymphomatous polyposis of gastrointestinal tract, report of, 247-256
of stomach and duodenum, metastases to, from carcinoma of breast, adrenal steroid-induced, 426-431
Succinylsulfathiazole (Sulfasuxidine), effect of, on radiation response of canine small intestine, report of, 862-874
SUCKOW, EARL E., *jt. au.*, 1205-1215
SUGAR, MAX, and WATKINS, CHARLES: Some observations about patients with a breast mass, 979-988
Sulfasuxidine (succinylsulfathiazole), effect of, on radiation response of canine small intestine, report of, 862-874
Sulphydryl (SH), compounds, effect of, on radiation sensitivity of mouse tumors, report of, 118, 119-120
SULLIVAN, ROBERT D., *jt. au.*, 744-752
SULLZEFF, V., *jt. au.*, 344-352, 1175-1182
Surgery, *See also names of specific operations*
as therapy for cancer of vulva, report of, 13-20
for giant follicular lymphoblastoma, report of, 772-773, 774
for multiple oral carcinoma, use of, 515
combined with chemotherapy, in treatment of malignant tumors, results of, 1197-1204

Surgery—(Contd.)

- (cruciate incision), effect of, on radiation sensitivity of mouse tumors, report of, 123-124
for squamous cell carcinoma of esophagus, stromal inflammatory cell infiltration as prognostic factor in evaluating effectiveness of, 921-933
in treatment of nonbenign bronchial carcinoids and cylindromas, results of, 554-546
of postmastectomy lymphangiosarcoma developing in lymphedematous arm, role of, 210-212
radical, attempted palliation of pelvic and abdominal carcinomatosis primary in ovaries by, 384-388
tumor cells disseminated during, effectiveness of small amounts of preoperative irradiation in preventing growth of, 284-295
Survival, in cases of adenoid cystic carcinoma, results of treatment and, 1247-1250
of patients with lung cancer treated by pulmonary resections, relation of certain factors to, 1251-1258
rates of, in cases of cancer of vulva, report of, 15-20
Syphilis, as etiologic factor in cancer of esophagus, 391, 397, 398, 401
TAKAHASHI, KIICHI: Squamous cell carcinoma of the esophagus. Stromal inflammatory cell infiltration as a prognostic factor, 921-933
TALLEY, R. W., *jt. au.*, 131-152, 493-495
TAN, EDWARD, *jt. au.*, 1259-1265
TANIGUCHI, LILIAN, *jt. au.*, 1074-1080
TAYLOR, SAMUEL G., III, *jt. au.*, 644-648
TDC. *See* "True diagnostic gap"
TELLEM, MILTON; STAHL, THEODORE, and MERANZE, DAVID R.: Carcinoma of the thyroid gland: an explanation for its varying incidence in surgically removed nodular thyroid glands, 67-72
TEM. *See* Triethylene melamine
Temperature, effect of, on radiation sensitivity of mouse tumors, 121-123
TEPA. *See* Triethylenephosphoramide
Teratocarcinoma(s), in vitro and in vivo study of differentiation in, 1017-1029
development of cell lines from embryoid bodies and teratocarcinomas in vitro in, 1027
transplantation of cells (glass-adapted) from, in mice in, 1027
of embryoid bodies from explanted teratocarcinomas in, 1022-1024
of embryoid bodies from explants of embryoid bodies in vitro in, 1027
differentiation of explants of embryoid bodies grafted subcutaneously in animals in, 1025-1027
incidence of various tissues found in, 1025-1027
microscopic structure of tumors in, 1025
dominance of muscle tissue in, 1025-1027
transplantation of teratocarcinomas to mice in, 1027
embryoid bodies cultured in vitro in, 1024-1025
fixing of explants in, 1019
LS402A VII tumor used in, 1017
media fed to cultures in, 1017
to teratocarcinomas in, 1017
patterns established in cultured teratocarcinomas in, 1019
size of embryoid bodies in, 1017
of fragments of teratocarcinomas to be used for tissue culture in, 1017
use of Hank's balanced salt solution in, 1017
solid precursor stages of cystic embryoid bodies chosen in, 1017
type of development in vitro in, 1019-1022
process of simplification in, 1019-1022
well differentiated polycystic teratocarcinomas as donor tumors in, 1018

Teratocarcinoma(s)—(Contd.)

- amount of differentiated somatic structures present in, 1018
commonly seen tissues in, 1018
embryonal carcinoma cells present in, 1018
Testis(es), bilateral carcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 243
malignant lymphomas of, in African children, pathological entity of, 275
spermatocytic seminoma of, 788-794
case reports, 788-790
clinical features in, 793
comparative morphology in, 793
comparison of classical seminoma and spermatocytic seminoma in, 791
microscopy of tumor in, 788-790
appearance of nuclei in, 789-793
size of cell in, 789-793
pathological findings in, 791-793
postoperative radiation used in, 788, 790
resemblance to canine seminoma in, 793, 794
Testosterone, effect of, on radiation sensitivity of mouse tumors, 119, 120
Therapy, for metastatic breast cancer, effect of androstetrone on clinical course and hormonal excretion in, 195-198
palliative, with deep X rays, for multiple polyposis of gastrointestinal tract, report of, 256
prophylactic, of plantar and palmar nevi, dissenting view on, 102-104
THOMAS, LOUIS B.; FORKNER, CLAUDE E., JR.; FARR, EMIL, III; BESSA, BYRON E., JR., and STABENAU, JAMES R.: The skeletal lesions of acute leukemia, 608-621
Thrombus(i), multiple, and carcinoma of pancreas, morphological study of, 944-952
Thymidine, tritiated, in labeling tissues in vitro, use in comparing rates of cell proliferation in duct epithelium, fibroadenoma, and carcinoma of breast, report of, 639-643
Thyroid, carcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 244
in childhood, report of, 734-743
clinical data in, 736-737
disease stage and survival in, 740-742
follow-up data in, 739-741
methods and results of treatment for patients followed more than 10 years in, 739, 740
staging of tumors and, 740-741
suspected hazards of pregnancy and, 741
geographical location in, 734-735
mortality data in, 741-742
pathological findings in, 738-739
bilateral involvement of thyroid lobes in, 739
change in tumor type in, 739
follicular cancer in, 738-739
papillary cancer in, 738-739
undifferentiated cancer in, 738-739
medullary as, 739
small cell as, 739
spindle and giant cell as, 739
previous irradiation in, 735-736
site of disease and survival in, 740-742
therapy in, 737-738
varying incidence of, in surgically removed nodular thyroids, explanation for, 67-72
approach to problem of, 67
comparison of clinically diagnosed solitary nodular goiter and pathologically diagnosed multinodular lesions in, 70-71
criteria for preoperative selectivity in, 68-69
clinical impression alone and, 68
patient age and sex alone and, 68, 69
scintiscanning alone and, 68-69
use of radioactive iodine

Thyroid, carcinomas of—(Contd.)

- (181) in, 68, 69
for carcinoma diagnosis in, 69-70
incidence of carcinoma based on clinically diagnosed nodular goiter in, 68
of carcinoma based on histological findings for operative specimens in, 68
operative procedures in, 68
preoperative classification of patients studied in, 67-68
involvement of, by giant follicular lymphoblastoma, report of, 758-759
malignant lymphomas of, in African children, pathological entity of, 275
Tissue(s). See also specific names
cultures, used in confirmation of epithelial origin of giant cell carcinoma of lungs, report of, 1052-1056
external, soft, as location of myxoma in children, report of, 633-634
from duct epithelium, fibroadenoma, and carcinoma of breast, labeled in vitro with tritiated thymidine, comparison of rates of cell proliferation of, 639-643
human, hydrolysis of naphthylchloroacetamide by, implications for chemotherapy of cancer, 296-299
malignant and benign neoplastic, cytochemical study of esterase activity of, 369-377
mouse, hydrolysis of naphthylchloroacetamide by, implications for chemotherapy of cancer, 296-299
soft, multiple congenital neoplasms of, report of, 913-920
appearance of sections of lesions in, 914, 915-917
stains used in, 917
approximate number of lesions found in, 915
case reports, 913-914
deposits of hemosiderin seen in some cases of, 913, 918, 919
difficulty in distinguishing benign and malignant lesions in, 917-919
distribution of lesions in, 915
etiology of, 919
familial relationship of individuals described in, 913
past history of cancer among relatives of family described in, 913
pathological findings in, 914, 915-917
prognosis in, 919-920
degree of differentiation and cellularity and age of patient in, 920
size of lesions in, 915, 918
well differentiated pattern of tumors seen in some areas of, 919
Tobacco. See also Cigarette(s) and Smoking
as etiological factor in cancer of esophagus, study of, 389-413
microscopic properties of basement membrane and elastic fibers of trachea and bronchus in relation to, 1175-1182
role of acidic fractions of, as promoters in carcinogenesis, study of, 1306-1315
biological results in, 1309-1310, 1312
promoting effect of phenolic fraction found in, 1309-1310
studies in, 1308
application method used in, 1308
amounts of material applied to mice in, 1308
breakdown of materials, 1308
division of study into series in, 1308
method of testing promoting activity of cigarette smoke condensate and its acidic fractions in, 1308
use of Swiss (Millerton) female mice in, 1308
chemical results in, 1307, 1308-1309
fractionation of phenolic fraction in, 1307

Tobacco, role of acidic fractions of—(Contd.)

- procedure for separation of tobacco smoke condensate into fractions in, 1307
nature of promoters in, 1311
role of initiators in, 1311-1312
results of short term tests in, 1310
test substances used in, 1307-1308
chromatography of 7,12-dimethylbenz[a]anthracene (DMBA) on Florisil in, 1307
preparation of test phenol in, 1307
purification of analytic reagent acetone in, 1308
of benz[a]pyrene by chromatography on alumina in, 1307
types of cigarettes used in, 1307
American nonfiltered king-sized cigarettes as, 1307
"spinnach" cigarettes as, 1307
TOKER, CYRIL: Some observations on Paget's disease of the nipple, 653-672
TOLLES, W. E.; HOWARD, W. J., and BOSTROM, R. C.: A study of the quantitative characteristics of exfoliated cells from the female genital tract. I. Measurement methods and results, 437-454
HOWARD, W. J., and BOSTROM, R. C.: A study of the quantitative characteristics of exfoliated cells from the female genital tract. II. Suitability of quantitative cytological measurements for automatic prescreening, 455-468
Tongue, multiple carcinomas, 512-516
"Total diagnostic gap," in biomathematical approach to tumor growth, report of, 1272-1294
silent interval," in biomathematical approach to tumor growth, report of, 1272-1294
Toxicity, manifestations of, from 6-aminonicotinamide (6-AM) therapy for disseminated cancer, 646, 647
of benzyl-N-bis(ethylenimido)phosphorocarbamate (AB-103), report of, 856-858, 860-861
of chlorambucil, in treatment of childhood cancer, incidence of, 39, 41
resulting from use of 5-fluorouracil (5-FU), clinical evaluation of, 132-136, 140-145
classification of, 132-133
complications of, 133-135
causes of death from, 133-134
manifestations of, 132, 133
changes in myeloid-erythroid ratio compared with degree of leukopenia as, 133, 134
Trachea, and bronchus, of smokers and nonsmokers, microscopic properties of basement membrane and elastic fibers of, 1175-1182
classification of smoking habits used in, 1175
correlation of type of epithelium, basement membrane, and smoking habits in, 1181
description of morphological classes in, 1176-1177
enzymatic digestion techniques used in, 1175-1176
fixing of tissue specimens in, 1175
use of Bouin's fluid in, 1175
of Carnoy's fluid in, 1175
method of measuring thickness and distribution of basement membrane in, 1176
areas measured in, 1176
use of ocular micrometer in, 1176
role of edema in connective tissue adjacent to basement membrane in, 1177
results for elastic fibers of trachea and bronchus in, 1178-1179
of measurements in, 1176, 1177-1178, 1179-1181
average thickness of membrane in, 1178
significance of differences found in, 1178
type of epithelium, basement membrane, and patient smoking habits in, 1178, 1179
using periodic acid-Schiff stains in, 1179-1180
staining properties seen in, 1177

Trachea, and bronchus—(Contd.)

- type of epithelium and average thickness of basement membrane in, 1179
Tracheitis sicca, use of pancreatic deoxyribonuclease (dornase) in treatment of, 416, 418, 420
TRAAMS, E. G., Jr. *et al.*, 953-956
Transaminase, glutamic oxalacetic (GOT), levels of, in serum of patients with carcinoma of breast, 726-733
glutamic pyruvic (GPT), levels of, in serum of patients with carcinoma of breast, report of, 726-733
Transplant(s), cancer, human, during experimental chemotherapy, growth curves of, 1117-1121
amount and type of necrosis in, 1118
comparison of control and tumor animals in, 1118
composite growth curves for tumors obtained in, 1118-1119
effect of amethopterin (4-amino-N¹⁰-methyl pteroylglutamic acid, Methotrexate) on growth rates in, 1118-1121
doses used and, 1118, 1119-1121
phases of growth in, 1119
everting of cheek pouch containing tumor in, 1117
formulas for calculation of tumor volume in, 1118
lag phase for Me 1 tumor growth curve and, 1119
for Pitt 41 tumor growth curve and, 1119
log phase for Me 1 tumor growth curve and, 1119, 1120
for Pitt 41 tumor growth curve and, 1119, 1120
method of transplanting tumor in, 1117
relationship between tumor volume and time after treatment of malignant melanoma (Me 1) with Methotrexate in, 1120
between tumor volume and time after treatment of thyroid cancer (Pitt 41) with Methotrexate in, 1120
study of tissue reaction around tumor transplant in, 1117-1118
subcutaneous injections of cortisone (Cortone Acetate) used in, 1117
time of tumor measurements in, 1118
use of golden hamsters (*Mesocricetus auratus*) in study of, 1117
of human malignant melanoma (Me 1) in study of, 1117
of human thyroid cancer (Pitt 41) in study of, 1117
Transplantation, of human cancer, growth curves of, during experimental chemotherapy, report of, 1117-1121
tumor, in mice, effect of growth hormone and adrenocorticotrophic hormone on, 507-511
use in experimental study of effectiveness of small amounts of preoperative irradiation in preventing growth of tumor cells disseminated at surgery, 284-295
Trichophytosis, skin test reactions to, in cases of Hodgkin's disease and lymphosarcoma, report of, 597-607
Triethylene melamine (TEM), in therapy of giant follicular lymphoblastoma, report of, 771, 772-773
radioactive-labeled (C¹⁴-labeled TEM), excretion and distribution studies in cancer patients with, 953-956
Triethylene thiophosphoramide (TSPA), comparison of, with preoperative irradiation, in prevention of growth of tumor cells disseminated at surgery, report on, 292, 293
Triethylene thiophosphoramide (TSPA), radioactive-labeled (P³²-labeled TSPA), excretion and distribution studies in cancer patients with, 953-956
Triosephosphate isomerase, activity of, in human blood serum from normal individuals and from individuals with pathological conditions, 1166-1174
Tritium (H³), labeling of cells from nutritional cirrhosis in autoradiographs with, 963-970
"True diagnostic gap" (TDO), in biomathematical approach to tumor growth, report of, 1272-1294
or "total survival time," in bio-

"True diagnostic gap"—(Contd.)

- mathematical approach to tumor growth, report of, 1272-1294
- silent interval (tsi), in biomathematical approach to tumor growth, report of, 1272-1294
- Tsi. See "True silent interval"
- TSPA. See Triethylene thiophosphoramide
- Tst. See "True—" or "Total survival time"
- Tuberculin (PPD), skin test reaction to, in cases of Hodgkin's disease and lymphosarcoma, report of, 597-607
- Tuberculosis(es), as possible etiological factor in peripheral lung cancer arising in scars, report of, 1213
- Tubule(s), mesonephric, derivation of tumors of female genital tract from, 711-725
- TUCKER, J. L., JR., and TALLEY, R. W.: Prolonged intraarterial chemotherapy for inoperable cancer: a technique, 493-495
- Tumor(s), adenocarcinomatous, of ovary and uterus, occurring as coexistent or sequential primary neoplasms, report of, 817-826
- adenoid, cystic, clinicopathological study of, 1235-1250
- of bronchus, report of, 161-178
- ameloblastic, adenomatoid, in man, classification, histopathology, and clinical behavior of, 81-82
- comparison of with ameloblastoma, 76
- aortic body, comparative study of, 779-787
- calcifying, epithelial, in man, classification, histopathology, and clinical behavior of, 80-81
- carcinoid, of bronchus, 161-178
- of small intestine, functioning, data for, 908-910
- carcinosarcoma 256 Walker as, use of "aging" in reducing development of, 1057-1059
- age of tumor cell suspension and delay in tumor development in rats after injection in, 1058-1059
- development of tumors in, 1057-1059
- method of investigation in, 1057
- determination of tumor cells in suspension in, 1057
- eosin solution used in staining in, 1057
- maintenance of solutions in, 1057
- preparation of tumor cell suspensions in, 1057
- repetition of experiments in, 1057
- Tyrod's solution used as diluent in, 1057
- use of Hanks's balanced salt solution in, 1057
- of Sprague-Dawley female albino rats in, 1057
- carotid body, study of, 779-787
- cells, carcinosarcoma 256 Walker, in rats, influence of thermal stress and changes in body temperature on, 111-116
- disseminated at surgery, effectiveness of small amounts of preoperative irradiation in preventing growth of, experimental study of, 284-295
- amount of radiation exposure used in, 286, 288-294
- effects on growth rate of primary tumors after irradiation in vivo in, 287, 288
- method of determining end points for different routes of injection in, 286
- for intramuscular and axillary routes, 286-287
- for intravascular route, 286
- of injecting in, 285-286
- of irradiating transplanted primary tumor in, 286
- method of measuring primary tumor size in, 287
- of preparing inoculum in, 285
- number of "takes" after different amounts of radiation exposure in, 289, 290, 291
- quantity and dilution of suspension injected intravenously in, 286

Tumor(s), ameloblastic—(Contd.)

- results of ancillary studies in, 292
- effect of fractionated radiation doses in, 292, 293
- of radiomimetic drug (triethylene thiophosphoramide, TSPA) in, 292, 293
- of disseminating in vivo irradiated cell suspension intramuscularly and into axillary wound in, 288-292
- effect of altering number of cells injected in, 289
- various dilutions of cell suspensions used in, 289, 290, 291
- of intravenous injection of cell suspensions from in vivo irradiated tumor in, 288, 289, 292
- time period between radiation of primary tumor and artificial dissemination of cells in, 293
- transplanted mouse tumors in, 285
- carcinoma dba59 as, 285
- lymphoma CD5 as, 285
- melanoma S91 Cloudman as, 285
- sarcoma dba49 as, 285
- sarcoma T241 Lewis as, 285
- epidermoid cellular structure of, and incidence of cancer in families of patients with cervical cancer, 183, 186, 193
- epithelial, role of hair follicle in origin and evolution of, 801-816
- basic tumor types and mode of evolution in, 803-809
- bud-shaped keratoacanthoma (type 1) in, 804-806
- carcinogenesis in glabrous epithelium in, 813
- dome-shaped and berry-shaped keratoacanthomas (types 2 and 3) in, 804, 807-809, 812
- location of, 807-809
- pseudocarcinomatous infiltration in, 809
- hair follicle as repository for carcinogens in, 811, 813
- growth cycle and carcinogenesis in, 813-815
- effect of hair regeneration in, 814-815
- experimental studies in, 813-815
- papillomas in, 809, 812, 814
- role of hair growth cycle and "hair follicular cycle" in, 802-803
- morphological changes that occur during, 802, 803
- active phase ("anagen") in, 802-803
- resting phase ("telogen") in, 802-803
- sebaceous adenomas in, 809
- significance of biological behavior of benign epithelial tumors in, 809-811
- single cell or multiple cell origin of tumors in, 815
- glomus, in children, 555-566
- age, sex, and race in, 556-564
- case reports, 561-563
- clinical characteristics in, 559-560
- familial aspects in, 556-561
- follow-up in, 557-563
- incidence of multiple tumors in, 556-564
- infiltration in, 559, 560-563, 564
- location of tumors in, 556-564
- malignant nature of tumors in, 563
- microscopic features in, 559
- main histological variants in, 559
- recurrences in, 557-563
- relationship to hemangiopericytomas in, 563-564
- treatment in, 557-564
- tumor size in, 557-563
- growth of, biomathematical approach to, 1272-1294
- differentiation of primary tumors from multiple metastases or from satellite lesions in, 1289-1293
- mathematical basis for evaluation

Tumor(s)—(Contd.)

- of exponential tumor growth in, 1272-1276
- definition of "silent interval" in, 1274-1275
- of "true silent interval" (tsi) 1274-1275
- derivative tumor growth concepts employed in, 1273
- equations derived in, 1273
- methodological considerations in, 1276-1278
- calculation of time of initial tumor implantation in, 1277
- classes of patients in, 1277-1278
- criteria for selection of patients in, 1276
- justifications for use of specific roentgenographic diameters as measures of tumor volume in, 1277
- method of measuring tumor volume in, 1276-1277
- equation for determining tumor volume in, 1277
- problem of biological predetermination in, 1291
- of cancer cell dormancy in, 1289-1291
- of determination of total tumor duration in, 1292-1293
- results for primary pulmonary neoplasms in, 1278-1286
- age, race, and sex of patients in, 1278-1281
- confirmation of diagnoses in, 1278
- diagnostic gaps recorded in, 1280-1286
- exponential growth curves found in, 1278-1286
- findings on posterior-anterior radiographs in, 1279-1286
- prediction of post-therapeutic tumor regression in, 1281-1286
- 6-diazo-5-oxo-L-norleucine (don) therapy and, 1281, 1282
- radiation therapy and, 1281
- tumor-doubling times found in, 1279-1286
- for secondary pulmonary neoplasms in, 1286-1289
- age, race, and sex of patients in, 1286, 1287
- direct measurement data for, 1286, 1287
- effect of radiation in, 1288
- exponential growth curves found in, 1287
- tumor size correlations used in, 1273-1274
- linear scale showing size-time correlations in, 1274
- use of diagnostic and survival intervals in, 1275-1276
- "apparent diagnostic gap" (ADG) and, 1275
- "apparent survival time" (AST) and, 1275
- "total diagnostic gap" and, 1275
- "total silent interval" and, 1275
- "true diagnostic gap" (TDG) and, 1275
- "true—" or "total survival time" (TST) and, 1275-1276
- "true silent interval" (TSI) and, 1275
- validity of growth rate studies in, 1289-1294
- histiocytic, in children, report of, 469-482
- in infants and children, malignant hemangioendothelioma as, report of, 1186-1196
- jugal body, study of, 779-787
- larynx, primary, incidence of lymph node metastases from and their relationship to certain characteristics of primary tumor, 55-66
- lung, in mice, after exposure to cigarette tar and beta radiation, incidence of, 346-347
- malignant, combined surgery and chemotherapy for, 1197-1204
- clinical trials in, 1199
- selection of cases of bronchogenic carcinoma for study in, 1199
- structure of drugs used in, 1199

Tumor(s), malignant—(Contd.)

- dose schedules employed in, 1198
effect on circulating cancer cells in, 1198
experimental animal groups in, 1197, 1198-1199
incidence of alopecia in, 1200
incidence of leukopenia in, 1200-1201
treatment of, with prednisone (Ultracorten), in, 1200
mortality rate in, 1200-1201
previous experimental studies of, 1197-1199
using actinomycin C (Sanamycin) in rats in, 1197
using ascites tumor Ehrlich cells in, 1197
using 8-azaguanine (Azan) in rats in, 1197
using carcinosarcoma 256 Walker tumor cells in, 1197
using E-39 (2,5-bis(ethylmimino)-3,6-dipropoxy-1,4-benzoxquinone) in rats in, 1197
using Mitomen (Nitromin, 2,2-dichloro-N-methyl-N-oxide hydrochloride) in rats in, 1197
using reticulum cell-like Yoshida tumor cells in, 1197
regimens used in, 1199-1200
taking of blood counts in, 1200
multiple, primary, different tissues or organs of origin, 231-237
frequency of, in cases of carcinoma tumors of small intestine, report of, 902-903, 905
of multicentric origin, 238-248
over-all report of, 221-230
occurrence of, with Bowen's type of intraepithelial and infiltrative carcinoma of vulva, incidence of, 320-321, 328-329
predisposition for organs of female genital tract in, 320-321, 329
mammary gland, in mice, after exposure to cigarette tar and beta radiation, incidence of, 347, 348, 350
melanotic, of infancy, appearing in adult, report of, 1090-1108
mesonephric, of female genital tract, report of, 711-725
mouse, effect of certain chemical and physical agents on radiation sensitivity of, 117-125
criteria for remissions in, 118-119
effect of incision in, 123-124
percentage of tumor remissions in control groups in, 118
protective effect of sulphydryl (SH) compounds in, 118, 119-120
comparison with effect of anoxia alone in, 122-123
tumor regression and, 123
effect of cold and anoxia in, 122-123
of heat in, 121
simultaneous radiation and, 121
damage to normal tissues resulting from, 121
method of plotting tumor regression in, 121
effect of hormones in, 120-121
use of cysteine as, 120
results of, 118, 119
use of glutathione as, 120
results of, 118, 119
results with testosterone as, 119, 120
comparison with glutathione in, 120
role of radiation and, 119, 120
radiation response in, 118-119
technique and dose used in, 117-118
determination of optimum dose range in, 118
results obtained in, 119-124
spontaneous mammary adenocarcinoma used in, 117
tumor size in, 117
use of DBA mice in, 117
mucocutaneous, of bronchus, report of, 168, 169, 175
multiple, congenital, of soft tissues, report of, 913-920

Tumor(s)—(Contd.)

- myxomatous, in children, 629-635
nerve trunk, in rabbits, study of, 1030-1035
odontogenic, in man and domesticated animals, classification, histopathology, and clinical behavior of, 73-101
comparison of ameloblastoma and craniopharyngioma in, 74
complex odontoma as, 88-89
clinical behavior of, 89
compound odontoma as, 89-90
microscopic appearance of, 89
epithelial odontogenic tumors with induction in, 82-90
ameloblastic fibroma as, 82-85
ameloblastic odontoma as, 87-88
ameloblastic sarcoma as, 85-86
dentinoma as, 86-87
odontoma as, 87
epithelial odontogenic tumors without induction in, 76-80
ameloblastic adenomatoid tumor as, 81-82
comparison of, with ameloblastoma, in, 76
microscopic appearance of, 82
theory of origin of, 82
ameloblastoma as, 76-80
comparison of, with ameloblastic adenomatoid tumor in, 76, 81
division of, into solid and cystic types, in, 77
predominant patterns in, 77-78
calcifying epithelial odontogenic tumor as, 80-81
odontogenic tumors of domesticated animals in, 95-96
odontogenic tumors of mesodermal origin in, 89, 90-95
cementifying fibroma as, 90-95
gigantiform or monstrous cementoma as one of, 93-95
odontogenic myxoma and odontogenic fibroma as, 89, 90
potentiality of oral epithelium in, 74, 75
tabulation of epithelial and mesodermal types in, 74
of reticuloendothelial or lymphoreticular system, in African children, clinical syndrome of, 258-269
ovarian, in mice, after exposure to cigarette tar and beta radiation, incidence of, 348-349, 351
"primary," spontaneous development of metastases after removal of, standardization protocol of 5 animal tumors used in study of, 875-882
pseudosarcomatous facitis as, in children, report of, 1216-1222
pulmonary, in newborn mice after exposure to carcinogenic chemicals, report of, 308-317
association of with malignant lymphoma in, 308-317
retinal anlage, congenital melanocarcinoma, melanotic adamantinoma, progonoma, and pigmented epulis of infancy, report of case in adult, 1090-1108
spermatocytic seminoma of testis as, report of, 788-794
transplantation of suspensions of, in patients with advanced cancer, quantitative studies of, 971-978
transplanted, effect of growth hormone (GH) and adrenocorticotrophic hormone (ACTH) on, 507-511
beef growth hormone as inactivated GH in, 507
determination of carcass weight change after primary tumor dissection in, 507-508
of size categories for pulmonary tumors in, 507
dye-staining for lungs in, 507
effect of nutritional factors in, 510
on nitrogen balance in, 510
measurements per animal obtained at end of experiments in, 508-511
differences in growth in, 508-511
in rate of formation of me-

Tumor(s), transplanted—(Contd.)

- metastases in, 508-511
rate of tumor take in, 509, 511
Somar-A (bovine) as GH in, 507
subcutaneous injection of hormones in, 507
Cortrophin-Zinc as ACTH in, 507
use of DBA/p gray mice in, 507
of sarcoma dba49 in, 507
uterine, leiomyosarcoma in situ and, 483-492
vascular, in infants and children, report of, 1186-1196
UHLIN, ALFRED, *jr.* *et. al.*, 841-852
ULM, AARON H., *et. al.*, 744-752
Upper alimentary tract. *See also specific names of organs*
cancer of, etiological factors in, 410
Upper respiratory tract, involvement of, by giant follicular lymphoblastoma, report of, 758
Urethan (ethyl carbamate), as carcinogenic chemical, in newborn mice, report of, 308-317
Urinary tract. *See also specific names of organs*
epitheliomas of, as multiple primary malignant tumors of multicentric origin, report of, 240-242
Urine, determination of levels of urokinase in, in patients with carcinoma, report of, 889-896
of cancer patients, excretion and distribution studies of radioactively-labeled alkylating drugs in, 953-956
Urokinase, excretion of, in patients with carcinoma, report of, 889-896
classification of patients in, 890
formula (expressed as specific gravity coefficient) used in, 889
method of quantitative determination used in, 889-891
of statistical evaluation in, 890
possible explanations for low urokinase levels in patients with far advanced cancer in, 895
urine collection in, 889-890
values found for carcinoma patients and normal individuals in, 890-893
statistical comparison of, 891-892
by carcinoma groups in, 891-892
Uterus. *See also Cervix*
and ovary, adenocarcinomas of, occurring as coexistent or sequential primary neoplasms, report of, 817-826
case reports, 818-823
criteria used in classification in, 823-824
microscopic appearance of tumors in, 818-823
operative procedures used in, 818-823
pathological findings in, 818-823
possible factor of hormonal disturbance in, 824
preparation of tissue sections in, 818
principal characteristics of, 818-825
clinical findings in, 817, 818-823
prognosis in, 818-825
staining technique used in, 818
leiomyosarcoma in situ of, 483-492
analysis of data in, 485-487
coexistence of in situ elements with initial invasive sarcoma in, 485
gross appearance of tumors in, 485, 487, 491
location of myomas in, 486-487
microscopic features found in, 484-485
atypia of cells in, 484-485
cytoplasm in, 485
results with different stains in, 485
patients' age range in, 485-486
possible transition stages in, 491-492
problems involved in study of, 483
prognosis in, 492
results of follow-up examinations in, 487
staining of histological slides in, 483
residual carcinoma in, microscopic

- Uterus, problems involved in study of—
(Contd.)
study of margins of cone biopsy specimen of cervix in predicting, 795-800
- Vaccination, with bacille Calmette Guérin (BCG), in study of delayed skin response in Hodgkin's disease and lymphosarcoma, report of, 597-607
- Vaccine, bovine, wart, treatment of laryngeal papillomas with, preliminary report of, 636-638
- Vagina, and cervix, in situ epidermoid carcinoma of, development of after radiotherapy of cervical cancer, report of, 353-360
- carcinomas of, as multiple primary malignant tumors of multicentric origin, report of, 242
- cells exfoliated from, measurement method used in study of quantitative characteristics of, 437-453
- cytology of, histiocytes and histiocytic reaction in, survey of risks of false positive diagnosis in, 1223-1225
- appearance of histiocytes in pathological conditions in, 1223
- description of histiocytes in, 1223
- difficulties of differential diagnosis in presence of histiocytic reaction in, 1223-1225
- appearance of cytoplasm in, 1224-1225
- of nuclei in, 1224
- intense cell reaction in, 1224
- number of mitoses in, 1224
- Papanicolaou classifications used in, 1224
- histiocytic reaction seen in, 1223-1225
- normal appearance of histiocytes and, 1223
- routine smears of, single trophoblastic cells as source of error in interpretation of, 1183-1185
- changes compatible with pregnancy seen in smears in, 1184
- classification of smears in, 1184
- differential characteristics in, 1184
- differentiation between "third type" cells and trophoblastic cells in, 1183, 1185
- follow-up biopsies in, 1184
- history of vaginal bleeding in patients included in study of, 1184
- types of cells in, 1183-1185
- suitability of quantitative cytological measurements for automatic pre-screening of cells exfoliated from, report of, 455-468
- VAITKEVICIUS, V. K.; BRENNAN, M. J.; BECKETT, V. L.; KELLY, J. E., and TALLEY, R. W.: Clinical evaluation of cancer chemotherapy with 5-fluorouracil, 131-152
- VAN RYMENANT, M., *jt. au.*, 1166-1174
- Vasculature, of juvenile melanoma, report of, 1001-1004
- Vein(s), hepatic, anatomical arrangements of, in relation to middle lobe lobectomy for liver cancer, 1296-1300
- VERNEY, E. L., *jt. au.*, 1017-1029
- Virus(es). *See also names of specific organisms*
as possible agent involved in recurrent laryngeal papillomas, report of, 636-638
- in relation to incidence of malignant lymphoma in African children, possible "priming" action of infection by, 282
- infections by, relationship of, to leukemia in mice, report of, 496-501
- SE polyoma, and host cell relationship, difference in, between tubular epithelium and stroma in mouse kidneys infected with, 567-576
- Viscera, abdominal, malignant lymphomas of, in African children, pathological entity of, 273-275
- VOUTRA, NEDA, and FORAKER, ALVAN G.: Relative maturation of surface epithelium and gland involvement of metaplasia, dysplasia, and intraepithelial carcinoma of the uterine cervix, 1043-1051
- Vulva, cancer of, report of, 13-20
- associated cancers and, 14
- classification of cases by clinical
- Vulva, cancer of—(Contd.)
staging in, 15-16
- 3- and 5-year survival rates and, 15-16
- contraindications for radiation therapy in, 17, 19
- delay in treatment of, 14
- histological findings in, 14-15
- lymph node involvement and, 14, 15, 18-20
- major postoperative complications involved in, 18-19
- modes of therapy for, 15-19
- 3- and 5-year survival rates in relation to, 16-18, 19-20
- patients in study of, 13-14
- age and menopausal status in, 13
- presenting symptoms in, 13-14
- radiation therapy in, 17-18
- relation of lymph node metastases to survival in, 18-19
- of type of therapy to clinical stage in, 17
- to survival in, 15
- size of lesion in, 16-20
- 3- and 5-year survival rates in relation to, 16-17, 19, 20
- carcinoma of, as multiple primary malignant tumors of multicentric origin, report of, 242
- infiltrative squamous cell (epidermoid), report of, 330-343
- anatomical areas included in, 330
- clinical features in, 331-332
- criteria for inclusion of cases in, 330
- cytological grading classification used in, 331
- desmoplastic response of stroma to invading carcinoma and survival in, 341
- division of patients into groups in, 333
- gross description of primary lesion in, 332
- histological features considered in, 330-331
- location of primary lesion in, 332
- morphological features of tumors in, 333-337, 341, 342
- areas of in situ change at margins in, 335
- fibrous reaction in, 335
- Grade I in, 333
- Grade II in, 333
- Grade III in, 333
- leukoplakic vulvitis in, 335, 341, 342
- metastatic carcinoma in lymph nodes in, 335
- multicentric carcinomatous areas in, 335
- multiple foci of carcinoma in, 340
- possibility of "host response" in, 341
- reasons for delay in diagnosis in, 337-338
- standard life table methods, to determine survival time, used in, 331
- survival data in, 335-337
- 1:1 ratio in, 335
- treatment group and probability of survival in, 337-341
- types of therapy used in, 332-333
- "diversified" therapy in, 333
- radical vulvectomy and bilateral inguinal lymph node dissection in, 333
- intraepithelial and infiltrative, Bowen's type, report of, 318-329
- age at onset of menses for patients in, 319
- clinical findings in, 319-321
- differentiation of Bowen's disease and Paget's disease in, 318
- incidence of other malignant neoplasms and, 320-321, 328-329
- initial clinical descriptions of lesions in, 319-321
- lack of leukoplakic vulvitis in, 321, 327
- menopausal status in, 319
- Vulva, carcinoma of—(Contd.)
pathological findings in, 321-323, 327-329
- associated condylomata acuminata with, 321
- foci of infiltrative squamous cell carcinoma in, 321
- histological changes in epithelial surfaces, epidermis and/or mucosa in, 320-321
- lack of metastases to inguinal lymph nodes in, 321-322
- multicentricity of lesions in, 321
- possible common underlying factor in development of other malignant neoplasms and, 328
- role of arsenic as ingested carcinogen in, 321, 328
- presenting symptoms in, 319
- reasons for delay in diagnosis in, 327
- by clinician, 325
- by patient, 327
- recurrences in, 323, 325, 328
- survival rates in, 324
- therapy prior to surgery in, 319, 321, 328
- treatment and survival in, 323-325
- radical vulvectomy and bilateral inguinal lymph node dissection in, 323
- vulvectomy or local excision of lesion in, 323-324
- varieties of intraepithelial squamous cell carcinoma in, 325
- Vulvectomy, complete, as treatment of Bowen's type of intraepithelial and infiltrative carcinoma of vulva, report of, 323-324, 328-329
- radical, and bilateral inguinal lymph node dissection, as treatment for infiltrative squamous cell (epidermoid) vulvar carcinoma, results of, 333, 341-343
- Vulvitis, leukoplakic, in infiltrative squamous cell (epidermoid) vulvar carcinoma, significance of, 335, 341, 342
- WADE-EVANS, T., and LANGLEY, F. A.: Mesonephric tumors of the female genital tract, 711-725
- WATKINS, CHARLES, *jt. au.*, 979-988
- WATSON, WILLIAM L., *jt. au.*, 539-546
- WEETH, J. B., *jt. au.*, 195-198, 707-710
- Weight, loss of, in patients with giant follicular lymphoblastoma, 763
- WEINBERG, THOMAS, *jt. au.*, 369-377
- WEISS, LEO, and INGRAM, MARGARET: Adenomatoid bronchial tumors. A consideration of the carcinoid tumors and the salivary tumors of the bronchial tree, 161-178
- WERK, EMILE E., JR., *jt. au.*, 105-110
- WEST, MICHAEL, *jt. au.*, 726-733
- "Wet film" (s), in detection of carcinoma of lung, cytological use of, 1139-1143
- WEXLER, HILDA, *jt. au.*, 875-882
- WHITAKER, JOANNE, *jt. au.*, 1115-1116
- WILSON, FRANK C., JR., and ANDERSON, PHILIP C.: A dissenting view of the prophylactic removal of plantar and palmar nevi, 102-104
- WINDMILLER, JOAN, *jt. au.*, 1115-1116
- WINKELMANN, R. K.: Juvenile melanoma. A histochemical study of the enzymes in 2 cases, 1001-1004
- WINSHIP, THEODORE, and ROSVOLL, RANDI V.: Childhood thyroid cancer, 734-743
- WOOLF, CHARLES M.: The incidence of cancer in the spouses of stomach cancer patients, 199-200
- and ISAACSON, E. ARNOLD: An analysis of 5 "stomach cancer families" in the State of Utah, 1005-1016
- Wound(s), operative, significance of cancer cells found in, report of, 1041-1044
- WOYKE, STANISLAW: Experimental tumors of peripheral nerve trunks in rabbits, 1030-1035
- WUESTER, WILLIAM O., and KNAUER, WARREN H.: Complete right hepatic lobectomy for rare teratoid liver cancer in infancy, 361-368
- WYNDER, ERNEST L., and BROSS, IRWIN J.: A study of etiologic factors

- in cancer of the esophagus, 389-413
and HOFFMANN, DIETRICH: A study of tobacco carcinogenesis. VIII. The role of the acidic fractions as promoters, 1306-1315
- Xanthoma(s), fibrous, in children, report of, 469-482
case reports, 477
clinical features of, 473, 480
age at onset in, 473, 480
anatomical location of tumor in, 473, 480
average duration of tumor in, 473, 480
differential diagnosis in, 475-477
differentiation between histiocytomas and, 470
follow-up in, 477
pathological findings in, 473-474
after gross study in, 473, 480
size of tumor in, 473, 478
after microscopic examination in, 473-474
benign tumor in, 473-474
malignant variety of tumor in, 474
- Xanthoma(s), fibrous—(Contd.)
questionably malignant variety of tumor in, 474
- X irradiation. *See also* Radiation and Therapy
at body temperature below 10° C.,
cancericidal effects of, 1060-1062
effect of hypothermia on median lethal dose (LD₅₀) in, 1061
groups of rats in, 1060, 1061
basic control, 1060, 1061
hypothermic without X irradiation, 1060, 1061
X-irradiated hypothermic, 1060, 1061
X-irradiated normothermic, 1060, 1061
preparation of tumor suspension in, 1060
radiation factors involved in, 1060-1061
survival time in, 1060-1061
tumor size at death in, 1061
use of adult Lewis strain rats in, 1060
of Lewis lymphoma as tumor in, 1060
- X ray(s). *See also* Radiation and Therapy
therapy, after operation for primary Hodgkin's disease of lung, results of, 1151-1165
for postmastectomy lymphangiosarcoma developing in lymphedematous arm, report of, role of, 210-212
- YOKOO, HIDEJIRO, and SUCKOW, EARL E.: Peripheral lung cancers arising in scars, 1205-1215
- ZARAFONETIS, CHRIS J. D.: Reticulum cell sarcoma complicating atopic dermatitis. Prolonged remission following Potaba therapy, 5-12
- ZIMMERMAN, HYMAN J., *jr. et al.*, 726-733
Zona fasciculata, thickness of, in carcinoma of lung, report of, 107
glomerulosa, thickness of, in carcinoma of lung, report of, 107
reticularis, thickness of, in carcinoma of lung, report of, 107